

Prior Authorization Criteria

2025 MCSNP

Last Updated: 1/1/2025

ABIRATERONE

Products Affected

- Abiraterone Acetate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of prostate cancer. One of the following: 1) Disease is metastatic, 2) Disease is regional node positive (e.g., Any T, N1, M0), 3) Patient is in a very-high-risk group receiving external beam radiation therapy (EBRT), or 4) Positive pelvic persistence/recurrence after prostatectomy. Used in combination with prednisone or dexamethasone. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog [e.g. Lupron (leuprolide), Zoladex (goserelin), Trelstar (triptorelin), Vantas (histrelin), Firmagon (degarelix)] or 2) Patient received bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ABRYSVO

Products Affected

- Abrysvo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	All indications: Vaccine is being used for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV). Immunization in age 60 years of age and older: Patient has not received an RSV vaccine (i.e., Abrysvo, Arexvy) in their lifetime. Age greater than or equal to 60 years. Immunization in pregnancy: Patient has not received Abrysvo vaccine for the current pregnancy. Both of the following: 1) Will be used for active immunization of pregnant individuals at 32 through 36 weeks gestational age and 2) Will be used for the prevention of severe LRTD caused by RSV in infants from birth through 6 months of age.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

ADALIMUMAB-AATY

Products Affected

- Adalimumab-aaty 1-pen Kit INJ
80MG/0.8ML
- Adalimumab-aaty 2-pen Kit
- Adalimumab-aaty 2-syringe Kit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis (Dx) of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate (MTX), leflunomide, sulfasalazine. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Dx of moderately to severely active PJIA. Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. Psoriatic Arthritis (PsA) (Initial): Dx of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Plaque psoriasis (PsO) (Initial): Dx of moderate to severe chronic PSO. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week TF/C/I to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, OR calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Ankylosing Spondylitis (AS) (Initial): Dx of active AS. Minimum duration of a one-month TF/C/I to one NSAID (eg, ibuprofen, naproxen) at maximally tolerated doses. Crohn's Disease (CD) (Initial): Dx of moderately to severely active CD. One of the following: frequent diarrhea and abdominal pain, at least 10% weight loss, complications (eg, obstruction, fever, abdominal mass), abnormal lab values (eg, CRP), OR CD Activity Index (CDAI) greater than 220. TF/C/I to one of the following conventional therapies: 6-mercaptopurine (6-MP), azathioprine, corticosteroid (eg, prednisone), MTX. Uveitis (UV) (Initial): Dx of non-infectious uveitis classified as intermediate, posterior, or panuveitis.
Age Restrictions	N/A
Prescriber Restrictions	RA, AS, JIA: (Initial) Prescribed by or in consultation with a rheumatologist. PsA: (Initial) Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS: (Initial) Prescribed by or in consultation with a dermatologist. CD, UC: (Initial) Prescribed by or in consultation with a gastroenterologist. UV (initial): Prescribed by or in consultation with a rheumatologist or an ophthalmologist.

Coverage Duration	UC: (Initial) 12 wks. Other uses (initial): 6 months. All uses (reauth): plan year.
Other Criteria	<p>Ulcerative Colitis (UC) (Initial): Dx of moderately to severely active UC. One of the following: greater than 6 stools per day, frequent blood in the stools, frequent urgency, presence of ulcers, abnormal lab values (eg, hemoglobin, ESR, CRP), OR dependent on, or refractory to, corticosteroids. TF/C/I to one of the following conventional therapies: 6-MP, azathioprine, corticosteroid (eg, prednisone), aminosalicylate (eg, mesalamine, olsalazine, sulfasalazine). Hidradenitis suppurativa (HS) (Initial): Dx of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). RA, PJIA, PsA, AS, PsO, CD, HS, UV (Reauth): Patient demonstrates positive clinical response to therapy. UC (Reauth): For patients who initiated therapy within the past 12 weeks: Patient has clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on therapy for longer than 12 weeks: Patient demonstrates positive clinical response to therapy.</p>

ADALIMUMAB-ADBM

Products Affected

- Adalimumab-adbm
- Adalimumab-adbm Crohns/uc/hs Starter
- Adalimumab-adbm Psoriasis/uveitis Starter
- Adalimumab-adbm Starter Package For Crohns Disease/uc/hs
- Adalimumab-adbm Starter Package For Psoriasis/uveitis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis (Dx) of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate (MTX), leflunomide, sulfasalazine. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Dx of moderately to severely active PJIA. Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. Psoriatic Arthritis (PsA) (Initial): Dx of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Plaque psoriasis (PsO) (Initial): Dx of moderate to severe chronic PSO. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week TF/C/I to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, OR calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Ankylosing Spondylitis (AS) (Initial): Dx of active AS. Minimum duration of a one-month TF/C/I to one NSAID (eg, ibuprofen, naproxen) at maximally tolerated doses. Crohn's Disease (CD) (Initial): Dx of moderately to severely active CD. One of the following: frequent diarrhea and abdominal pain, at least 10% weight loss, complications (eg, obstruction, fever, abdominal mass), abnormal lab values (eg, CRP), OR CD Activity Index (CDAI) greater than 220. TF/C/I to one of the following conventional therapies: 6-mercaptopurine (6-MP), azathioprine, corticosteroid (eg, prednisone), MTX. Uveitis (UV) (Initial): Dx of non-infectious uveitis classified as intermediate, posterior, or panuveitis.
Age Restrictions	N/A
Prescriber Restrictions	RA, AS, JIA: (Initial) Prescribed by or in consultation with a rheumatologist. PsA: (Initial) Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS: (Initial) Prescribed by or in consultation with a dermatologist. CD, UC: (Initial) Prescribed by or in consultation with a gastroenterologist. UV (initial): Prescribed by or in consultation with a rheumatologist or an ophthalmologist.

Coverage Duration	UC: (Initial) 12 wks. Other uses (initial): 6 months. All uses (reauth): plan year.
Other Criteria	<p>Ulcerative Colitis (UC) (Initial): Dx of moderately to severely active UC. One of the following: greater than 6 stools per day, frequent blood in the stools, frequent urgency, presence of ulcers, abnormal lab values (eg, hemoglobin, ESR, CRP), OR dependent on, or refractory to, corticosteroids. TF/C/I to one of the following conventional therapies: 6-MP, azathioprine, corticosteroid (eg, prednisone), aminosalicylate (eg, mesalamine, olsalazine, sulfasalazine). Hidradenitis suppurativa (HS) (Initial): Dx of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). RA, PJA, PsA, AS, PsO, CD, HS, UV (Reauth): Patient demonstrates positive clinical response to therapy. UC (Reauth): For patients who initiated therapy within the past 12 weeks: Patient has clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on therapy for longer than 12 weeks: Patient demonstrates positive clinical response to therapy.</p>

ADEMPAS

Products Affected

- Adempas

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH. Chronic thromboembolic pulmonary hypertension (CTEPH): One of the following: A) Both of the following: 1) Diagnosis of inoperable or persistent/recurrent CTEPH and 2) CTEPH is symptomatic OR B) Patient is currently on any therapy for the diagnosis of CTEPH.
Age Restrictions	N/A
Prescriber Restrictions	PAH, CTEPH: Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH, CTEPH: plan year
Other Criteria	N/A

AIMOVIG

Products Affected

- Aimovig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Episodic Migraines (EM) (initial): Diagnosis of EM. Patient has greater than or equal to 4 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) Topiramate (Topamax), e) Venlafaxine (Effexor), f) Candesartan (Atacand). Chronic Migraines (CM) (initial): Diagnosis of CM. Patient has greater than or equal to 8 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) OnabotulinumtoxinA (Botox), e) Topiramate (Topamax), f) Venlafaxine (Effexor), g) Candesartan (Atacand). All Indications (initial): Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	EM, CM (initial, reauth): Plan year.
Other Criteria	EM, CM (reauth): Patient demonstrates positive clinical response to therapy. Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.

AKEEGA

Products Affected

- Akeega

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of prostate cancer. Disease is all of the following: a) metastatic, b) castration-resistant, and c) deleterious or suspected deleterious BRCA-mutated (BRCAm). Used in combination with prednisone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ALECENSA

Products Affected

- Alecensa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is anaplastic lymphoma kinase (ALK)-positive. One of the following: 1) Disease is one of the following: a) Recurrent, b) Advanced, or c) Metastatic, or 2) Used as adjuvant treatment following tumor resection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ALOSETRON

Products Affected

- Alosetron Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Irritable bowel syndrome (IBS) (Initial): Exclude if patient is of the male gender.
Required Medical Information	IBS (Initial): Diagnosis of chronic severe diarrhea-predominant IBS. IBS (Reauthorization): Symptoms of IBS continue to persist. Patient demonstrates positive clinical response to therapy.
Age Restrictions	IBS (Initial): 18 years and older.
Prescriber Restrictions	N/A
Coverage Duration	IBS (Initial): 12 weeks. IBS (Reauthorization): 6 months.
Other Criteria	IBS (initial): Trial and failure, contraindication, or intolerance to an anti-diarrheal agent [eg, loperamide].

ALPHA - 1 PROTEINASE INHIBITORS

Products Affected

- Aralast Np INJ 1000MG
- Prolastin-c INJ 1000MG/20ML
- Zemaira INJ 1000MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Alpha-1 antitrypsin (AAT) deficiency (initial): Diagnosis of congenital AAT deficiency. Diagnosis of emphysema. One of the following: Pi*ZZ, Pi*Z(null) or Pi*(null)(null) protein phenotypes (homozygous) or Other rare AAT disease genotypes associated with pre-treatment serum AAT level less than 11 µmol/L [eg, Pi(Malton, Malton), Pi(SZ)]. One of the following: 1) Circulating pre-treatment serum AAT level less than 11 µmol/L (which corresponds to less than 80 mg/dL if measured by radial immunodiffusion or less than 57 mg/dL if measured by nephelometry), or 2) Patient has a concomitant diagnosis of necrotizing panniculitis. Continued conventional treatment for emphysema (eg, bronchodilators).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	AAT deficiency (initial, reauth): plan year
Other Criteria	AAT deficiency (reauth): Patient demonstrates positive clinical response to therapy. Continued conventional treatment for emphysema (e.g., bronchodilators).

ALUNBRIG

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic, recurrent, or advanced NSCLC and tumor is anaplastic lymphoma kinase (ALK)-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

AMBRISENTAN

Products Affected

- Ambrisentan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH: Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: plan year
Other Criteria	N/A

APREPITANT

Products Affected

- Aprepitant CAPS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Chemotherapy-induced Nausea and Vomiting (CINV): Patient is currently receiving moderately or highly emetogenic chemotherapy. Patient is concurrently on both corticosteroid [eg, Decadron (dexamethasone)] and 5-HT3 receptor antagonist [eg, Aloxi (palonosetron), Anzemet (dolasetron), Kytril (granisetron), Zofran (ondansetron)]. Delayed Chemotherapy-induced Nausea and Vomiting Prevention: Patient is currently receiving highly emetogenic chemotherapy and corticosteroid [eg, Decadron (dexamethasone)], or patient is receiving an anthracycline [eg, Adriamycin (doxorubicin), Ellence (epirubicin)] and Cytosan (cyclophosphamide), or patient is currently receiving moderately emetogenic chemotherapy and was given aprepitant (oral or IV) on day 1 of chemotherapy. Postoperative Nausea and Vomiting (PONV): For the prevention of postoperative nausea and vomiting when administered prior to the induction of anesthesia.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Acute CINV, Delayed CINV: plan year. PONV: 1 month
Other Criteria	Subject to Part B vs. Part D review.

ARANESP

Products Affected

- Aranesp Albumin Free INJ
100MCG/0.5ML, 100MCG/ML,
10MCG/0.4ML, 150MCG/0.3ML,
200MCG/0.4ML, 200MCG/ML,
25MCG/0.42ML, 25MCG/ML,
300MCG/0.6ML, 40MCG/0.4ML,
40MCG/ML, 500MCG/ML,
60MCG/0.3ML, 60MCG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Anemia due to Chronic Kidney Disease (CKD) (Initial): Diagnosis of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) within 30 days of request. The rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. (Reauth): Diagnosis of CKD. Most recent or average (avg) Hct over 3 mo is 33% or less (Hgb 11 g/dL or less) for patients on dialysis, without ESRD OR Most recent or average (avg) Hct over 3 mo is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis OR Most recent or average (avg) Hct over 3 mo is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Documentation of a positive clinical response to therapy from pre-treatment level. Anemia w/ chemo (Initial): Other causes of anemia ruled out. Anemia w/ labs (Hct less than 30%, Hgb less than 10 g/dL) within prior 2 weeks of request. Cancer is non-myeloid malignancy. Patient is receiving chemo. (Reauth): Anemia by labs (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 weeks of request. Documentation of a positive clinical response to therapy from pre-treatment level. Patient is receiving chemo. Anemia in MDS (Init): Diagnosis of MDS. Serum erythropoietin 500 mU/mL or less, or transfusion dependent MDS. (Reauth): Most recent or avg Hct over 3 months was 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Documentation of a positive clinical response to therapy from pre-treatment level.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CKD(Init): 6 mo. CKD(reauth):plan yr. Chemo(init, reauth): 3 mo. MDS(init): 3 mo,(reauth): plan yr
Other Criteria	ESRD patients: Coverage is excluded under Medicare Part D for patients with ESRD on dialysis for any indication related or unrelated to treatment of ESRD since the payment for the drug is included in the ESRD PPS payment bundle. NON-ESRD PATIENTS for the following indications: Off-label uses (except Anemia in Myelodysplastic Syndrome (MDS):

	Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%.CKD (init, reauth), Chemo (init), MDS (init): Verify Fe eval for adequate Fe stores.
--	---

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cryopyrin-Associated Period Syndromes (CAPS): Diagnosis of CAPS, Familial Cold Autoinflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS). Deficiency of Interleukin-1 Receptor Antagonist (DIRA): Diagnosis of DIRA. Patient weighs at least 10 kg. Patient is currently in remission (e.g., no fever, skin rash, and bone pain/no radiological evidence of active bone lesions/C-reactive protein [CRP] less than 5 mg/L). Recurrent Pericarditis (initial): Diagnosis of recurrent pericarditis as evidenced by at least 2 episodes that occur a minimum of 4 to 6 weeks apart. Trial and failure, contraindication, or intolerance (TF/C/I) to at least one of the following: nonsteroidal anti-inflammatory drugs (e.g., ibuprofen, naproxen), colchicine, or corticosteroids (e.g., prednisone).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CAPS, DIRA: plan year. Recurrent Pericarditis (initial, reauth): plan year.
Other Criteria	Recurrent Pericarditis (reauth): Patient demonstrates positive clinical response to therapy.

AREXVY

Products Affected

- Arexvy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Immunization in age 60 years of age and older: Vaccine is being used for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV). Patient has not received an RSV vaccine (i.e., Abrysvo, Arexvy) in their lifetime. Age greater than or equal to 60 years. Immunization in age 50 through 59 years: Vaccine is being used for the prevention of LRTD caused by RSV. Patient has not received an RSV vaccine (i.e., Abrysvo, Arexvy) in their lifetime. Age 50 through 59 years. Patient is at increased risk for LRTD caused by RSV.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

ARIKAYCE

Products Affected

- Arikayce

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mycobacterium avium complex (MAC) lung disease: Diagnosis of Mycobacterium avium complex (MAC) lung disease. Used as part of a combination antibacterial drug regimen. Used in patients who do not achieve at least two negative sputum cultures after a minimum of 6 consecutive months of a multidrug background regimen therapy (e.g., a macrolide, a rifamycin, ethambutol, etc).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist.
Coverage Duration	Plan year
Other Criteria	N/A

ARMODAFINIL

Products Affected

- Armodafinil

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Obstructive sleep apnea (OSA) (Initial): Diagnosis (dx) of OSA defined by one of the following: a) 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or b) both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), AND 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work disorder (SWD) (Initial): Dx of SWD confirmed by one of the following: 1) Symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). Confirmation that no other medical conditions or medications are causing the symptoms of excessive sleepiness or insomnia. Narcolepsy (initial): Dx of narcolepsy as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	OSA, SWD: Initial, Reauth: 6 mo. Narcolepsy: Initial, Reauth: Plan Year
Other Criteria	OSA, Narcolepsy (Reauth): Patient demonstrates positive clinical response to therapy. SWD (Reauth): Patient demonstrates positive clinical response to therapy.

AUGTYRO

Products Affected

- Augtyro CAPS 40MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Disease is one of the following: a) advanced, or b) metastatic. Disease is ROS1-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

AUSTEDO

Products Affected

- Austedo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chorea associated with Huntington's disease: Diagnosis of Chorea associated with Huntington's disease. Tardive dyskinesia: Diagnosis of tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication or 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication.
Age Restrictions	N/A
Prescriber Restrictions	Huntington's disease chorea: Prescribed by or in consultation with a neurologist. Tardive dyskinesia: Prescribed by or in consultation with a neurologist or psychiatrist.
Coverage Duration	Plan year
Other Criteria	N/A

AYVAKIT

Products Affected

- Ayvakit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gastrointestinal stromal tumor (GIST): Diagnosis of GIST. One of the following: 1) Patient has unresectable, recurrent, or metastatic disease after failure on one approved therapy (e.g., imatinib, sunitinib, dasatinib, regorafenib, ripretinib) OR 2) Both of the following: a) Disease is one of the following: i) unresectable, ii) metastatic, iii) recurrent, iv) gross residual disease (R2 resection), v) residual disease with significant morbidity, vi) limited progression, or vii) resectable with significant morbidity AND b) Presence of platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations. Systemic Mastocytosis: Both of the following: 1) Diagnosis of one of the following: a) advanced systemic mastocytosis (AdvSM), b) aggressive systemic mastocytosis (ASM), c) systemic mastocytosis with an associated hematological neoplasm (SM-AHN), or d) mast cell leukemia (MCL) AND 2) platelet count is greater than $50 \times 10^9/L$. Ayvakit 25 mg - Indolent Systemic Mastocytosis (ISM): Diagnosis of ISM. Platelet count is greater than $50 \times 10^9/L$.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BALVERSA

Products Affected

- Balversa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Urothelial Carcinoma: Diagnosis of urothelial carcinoma (UC). One of the following: Locally advanced or metastatic. Presence of fibroblast growth factor receptor (FGFR) 3 genetic alterations. Disease has progressed on or after at least one line of prior systemic therapy (e.g., chemotherapy). One of the following: 1) Patient has received prior systemic therapy containing an immune checkpoint inhibitor (e.g., pembrolizumab, nivolumab, avelumab) or 2) Patient is not eligible for immune checkpoint inhibitor therapy (e.g., pembrolizumab, nivolumab, avelumab).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BENLYSTA

Products Affected

- Benlysta INJ 200MG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Systemic Lupus Erythematosus (SLE) (Initial): Diagnosis of active SLE. Autoantibody positive (ie, anti-nuclear antibody [ANA] titer greater than or equal to 1:80 or anti-dsDNA level greater than or equal to 30 IU/mL). Currently receiving at least one standard of care treatment for active SLE (eg, antimalarials [eg, Plaquenil (hydroxychloroquine)], corticosteroids [eg, prednisone], or immunosuppressants [eg, methotrexate, Imuran (azathioprine)]). Lupus Nephritis (Initial): Diagnosis of active lupus nephritis. Currently receiving standard of care treatment for active lupus nephritis (e.g., corticosteroids [e.g., prednisone] with mycophenolate or cyclophosphamide). SLE, Lupus Nephritis (Reauthorization): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	SLE, Lupus Nephritis (initial, reauth): 6 months
Other Criteria	N/A

BERINERT

Products Affected

- Berinert

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Treatment of hereditary angioedema (HAE) attacks (initial): Diagnosis of HAE. Diagnosis has been confirmed by C1 inhibitor (C1-INh) deficiency or dysfunction (Type I or II HAE) as documented by ONE of the following: a) C1-INH antigenic level below the lower limit of normal OR b) C1-INH functional level below the lower limit of normal. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
Age Restrictions	N/A
Prescriber Restrictions	HAE (initial): Prescribed by or in consultation with an immunologist or an allergist
Coverage Duration	Initial, Reauth: Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. Not used in combination with other approved treatments for acute HAE attacks

BESREMI

Products Affected

- Besremi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of polycythemia vera as confirmed by all of the following: 1) One of the following: a) Hemoglobin greater than 16.5 g/dL for men or hemoglobin greater than 16.0 g/dL for women, b) Hematocrit greater than 49% for men or hematocrit greater than 48% for women, or c) Increased red cell mass, AND 2) Bone marrow biopsy showing hypercellularity for age with trilineage growth (panmyelosis) including prominent erythroid, granulocytic and megakaryocytic proliferation with pleomorphic, mature megakaryocytes, AND 3) One of the following: a) Presence of JAK2 or JAK2 exon 12 mutation or b) Subnormal serum erythropoietin level. Both of the following: 1) Trial and failure, contraindication or intolerance (TF/C/I) to hydroxyurea, AND 2) TF/C/I to one interferon therapy (e.g., Intron A, Pegasys, etc.).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BEXAROTENE

Products Affected

- Bexarotene

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cutaneous T-cell lymphoma (CTCL): Diagnosis of cutaneous T-cell lymphoma (CTCL).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BEXSERO

Products Affected

- Bexsero

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroup B. Patient has not received or will not receive more than two Bexsero or Trumenba vaccines in the previous 12 months. One of the following: 1) Age lesser than or equal to 25 years, or 2) Patient is in a high risk group as defined by the Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year (2 vaccinations per year)
Other Criteria	N/A

BOSENTAN

Products Affected

- Bosentan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH AND PAH is symptomatic AND One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH
Age Restrictions	N/A
Prescriber Restrictions	PAH: Prescribed by or in consultation with a pulmonologist or cardiologist
Coverage Duration	PAH: Plan year
Other Criteria	N/A

BOSULIF

Products Affected

- Bosulif

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of chronic myelogenous/myeloid leukemia (CML).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BRAFTOVI

Products Affected

- Braftovi CAPS 75MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Patient is positive for BRAF V600 mutation. Used in combination with Mektovi (binimetinib). Colorectal Cancer: One of the following diagnoses: Colon Cancer or Rectal Cancer. One of the following: 1) Unresectable or advanced disease or 2) Metastatic disease. Patient has received prior therapy. Patient is positive for BRAF V600E mutation. Used in combination with one of the following: 1) Erbitux (cetuximab) or 2) Vectibix (panitumumab). Non-Small Cell Lung Cancer: Diagnosis of metastatic NSCLC. Patient is positive for BRAF V600E mutation. Used in combination with Mektovi (binimetinib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

BRIVIACT

Products Affected

- Briviact ORAL SOLN
- Briviact TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Partial-onset seizures: Diagnosis of partial-onset seizures.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BRONCHITOL

Products Affected

- Bronchitol

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic Fibrosis (CF) (initial): Diagnosis of CF. Patient has passed the Bronchitol Tolerance Test (BTT).
Age Restrictions	CF (initial): Patient is 18 years of age or older.
Prescriber Restrictions	CF (initial): Prescribed by or in consultation with a pulmonologist or specialist affiliated with a CF care center.
Coverage Duration	CF (initial): 6 months. CF (reauth): Plan year.
Other Criteria	CF (reauth): Patient demonstrates positive clinical response to therapy.

BRUKINSA

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mantle Cell Lymphoma (MCL): Diagnosis of MCL. Contraindication or intolerance to Calquence (acalabrutinib). Waldenstrom's Macroglobulinemia (WM)/Lymphoplasmacytic Lymphoma (LPL): Diagnosis of WM/LPL. Contraindication or intolerance to Imbruvica (ibrutinib). Marginal Zone Lymphoma (MZL): Diagnosis of MZL. Disease is relapsed, refractory, or progressive. Patient has received at least one anti-CD20-based regimen (e.g., rituximab, obinutuzumab). Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Contraindication or intolerance to Calquence (acalabrutinib). Follicular Lymphoma (FL): Diagnosis of FL. Disease is relapsed or refractory. Used in combination with Gazyva (obinutuzumab). Patient has received at least two prior lines of systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year.
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

BYDUREON

Products Affected

- Bydureon Bcise

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

BYETTA

Products Affected

- Byetta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

CABLIVI

Products Affected

- Cablivi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acquired thrombotic thrombocytopenic purpura (aTTP): Diagnosis of aTTP. First dose was/will be administered by a healthcare provider as a bolus intravenous injection. Used in combination with immunosuppressive therapy (e.g. rituximab, glucocorticoids). One of the following: 1) Used in combination with plasma exchange or 2) both of the following: patient has completed plasma exchange and less than 59 days have or will have elapsed beyond the last plasma exchange.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

CABOMETYX

Products Affected

- Cabometyx

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma (RCC): Diagnosis of advanced RCC. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. One of the following: a) Trial and failure, contraindication, or intolerance to Nexavar (sorafenib tosylate), or b) Both of the following: 1) Disease is Child-Pugh class A and 2) Patient has unresectable disease and is not a transplant candidate, or c) Both of the following: 1) Disease is Child-Pugh class A and 2) Patient has metastatic disease or extensive liver tumor burden, or d) Both of the following: 1) Disease is Child-Pugh class A and 2) Patient has liver-confined disease and is inoperable by performance status, comorbidity, or with minimal or uncertain extrahepatic disease. Differentiated Thyroid Cancer (DTC): Diagnosis of DTC. Disease is one of the following: a) locally advanced or b) metastatic. Disease has progressed following prior VEGFR-targeted therapy (e.g., Lenvima [lenvatinib], Nexavar [sorafenib]). Disease is radioactive iodine-refractory or ineligible.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CALQUENCE

Products Affected

- Calquence

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mantle Cell Lymphoma: Diagnosis of mantle cell lymphoma (MCL). Patient has received at least one prior therapy for MCL [e.g., Rituxan (rituximab)]. Chronic Lymphocytic Leukemia (CLL) or Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CAPLYTA

Products Affected

- Caplyta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Schizophrenia: Diagnosis of schizophrenia. Trial and failure, contraindication, or intolerance to two of the following: asenapine, aripiprazole, paliperidone, olanzapine, quetiapine (IR or ER), risperidone, or ziprasidone. Bipolar disorder: Diagnosis of bipolar I or II disorder (bipolar depression). Patient has depressive episodes associated with bipolar disorder. Used as monotherapy or as adjunctive therapy with lithium or valproate.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic Fibrosis (CF): Diagnosis of CF and lung infection with positive culture demonstrating Pseudomonas aeruginosa infection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

CHENODAL

Products Affected

- Chenodal

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of radiolucent gallstones. Patient has a well-opacifying gallbladder visualized by oral cholecystography. Trial and failure, contraindication or intolerance to ursodiol. Patient is not a candidate for surgery. Stones are not calcified (radiopaque) or radiolucent bile pigment stones.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, reauth: Plan year
Other Criteria	Reauth: Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment as evidenced by Oral cholecystograms or ultrasonograms.

CHOLBAM

Products Affected

- Cholbam

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Bile acid synthesis disorders due to single enzyme defects (BAS) (initial): diagnosis of a bile acid synthesis disorder due to a single enzyme defect. Peroxisomal disorders (PD) (initial): All of the following: 1) diagnosis of peroxisomal disorder, 2) patient exhibits at least one of the following: a) liver disease (eg, jaundice, elevated serum transaminases), OR b) steatorrhea, OR c) complications from decreased fat-soluble vitamin absorption (eg, poor growth), AND 3) Will be used as an adjunctive treatment.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by a hepatologist, medical geneticist, pediatric gastroenterologist, OR other specialist that treats inborn errors of metabolism.
Coverage Duration	Initial: 3 months Reauth: Plan year
Other Criteria	All uses (reauth): Patient demonstrates positive clinical response to therapy.

CLOBAZAM

Products Affected

- Clobazam

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Lennox-Gastaut syndrome: Diagnosis of Lennox-Gastaut syndrome. Used for the adjunctive treatment of seizures associated with Lennox-Gastaut syndrome. Dravet syndrome: Diagnosis of seizures associated with Dravet syndrome (DS). Used in combination with Diacomit.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CLONIDINE ER

Products Affected

- Clonidine Hydrochloride Er TB12

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of attention deficit hyperactivity disorder (ADHD).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

COMETRIQ

Products Affected

- Cometriq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Medullary Thyroid Cancer (MTC): Diagnosis of metastatic medullary thyroid cancer (MTC).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

COPIKTRA

Products Affected

- Copiktra

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Disease is relapsed or refractory. Trial and failure, contraindication, or intolerance to at least two prior therapies for CLL/SLL [e.g., Leukeran (chlorambucil), Gazyva (obinutuzumab), Arzerra (ofatumumab), Bendeka (bendamustine), Imbruvica (ibrutinib), etc.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CORLANOR

Products Affected

- Corlanor SOLN
- Ivabradine Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Chronic heart failure (CHF) (initial): Diagnosis of CHF. Patient has NYHA Class II, III, or IV symptoms. Patient has a left ventricular ejection fraction less than or equal to 35%. Patient is in sinus rhythm. Patient has a resting heart rate of greater than or equal to 70 beats per minute. Patient has been hospitalized for worsening HF in the previous 12 months. Trial and failure, contraindication, or intolerance to two of the following at a maximally tolerated dose: A) One of the following: 1) ACE inhibitor (e.g., captopril, enalapril, lisinopril), 2) ARB (e.g., candesartan, losartan, valsartan), or 3) ARNI (e.g., Entresto [sacubitril and valsartan]), B) One of the following: 1) bisoprolol, 2) carvedilol, or 3) metoprolol succinate extended release, C) Sodium-glucose co-transporter 2 (SGLT2) inhibitor [e.g., Jardiance (empagliflozin), Farxiga (dapagliflozin), Xigduo XR (dapagliflozin and metformin)], or D) Mineralocorticoid receptor antagonist (MRA) [e.g., eplerenone, spironolactone]. Dilated Cardiomyopathy (DCM) (initial): Diagnosis of heart failure due to DCM. Patient has NYHA Class II, III, or, IV symptoms. Patient is in sinus rhythm. Patient has an elevated heart rate. Trial and failure, contraindication or intolerance to one of the following: 1) Beta blocker (e.g., bisoprolol, metoprolol succinate extended release), 2) Angiotensin-converting enzyme (ACE) inhibitor (e.g., captopril, enalapril), or 3) Diuretic Agent (e.g., spironolactone, furosemide).</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CHF, DCM (Initial, reauth): plan year
Other Criteria	CHF, DCM (reauth): Patient demonstrates positive clinical response to therapy.

COSENTYX

Products Affected

- Cosentyx INJ 150MG/ML,
75MG/0.5ML
- Cosentyx Sensoready Pen
- Cosentyx Unoready

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Plaque psoriasis (Initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week trial and failure, contraindication, or intolerance (TF/C/I) to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. Minimum duration of a one-month TF/C/I to one non-steroidal anti-inflammatory drug (NSAID) (eg, ibuprofen, naproxen) at maximally tolerated doses. Non-radiographic axial spondyloarthritis (nr-axSpA, initial): Dx of active nr-axSpA with signs of inflammation. Minimum duration of a one-month TF/C/I to two non-steroidal anti-inflammatory drugs (NSAIDs) (eg, ibuprofen, naproxen) at maximally tolerated doses. Hidradenitis suppurativa (HS) (Initial): Diagnosis of moderate to severe HS.
Age Restrictions	N/A
Prescriber Restrictions	Plaque psoriasis, HS (Initial): Prescribed by or in consultation with a dermatologist. Psoriatic Arthritis (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. AS, nr-axSpA, ERA (initial): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	All uses (Initial): 6 months. All uses (Reauth): plan year.
Other Criteria	Enthesitis-Related Arthritis (ERA) (Initial): Diagnosis of active ERA. Minimum duration of a one-month TF/C/I to two NSAIDs (eg, ibuprofen, naproxen) at maximally tolerated doses. All indications (Reauth): Patient demonstrates positive clinical response to therapy.

COTELLIC

Products Affected

- Cotellic

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable or metastatic melanoma. Disease is positive for BRAF V600E or V600K mutation. Used in combination with Zelboraf (vemurafenib). Histiocytic Neoplasm: Diagnosis of one of the following: 1) Langerhans Cell Histiocytosis, 2) Erdheim-Chester Disease, or 3) Rosai-Dorfman Disease. One of the following: 1) Mitogen-activated protein (MAP) kinase pathway mutation, 2) No detectable mutation, or 3) Testing not available.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

CRINONE

Products Affected

- Crinone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	All indications: Excluded if for fertility uses.
Required Medical Information	Secondary amenorrhea: Diagnosis of secondary amenorrhea (the absence of menses in women who have already started menstruation who are not pregnant, breastfeeding, or in menopause).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CYCLOSET

Products Affected

- Cycloset

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diabetes Mellitus (Initial): Diagnosis of type 2 diabetes mellitus (DM). Both of the following: 1) Trial and failure, contraindication, or intolerance to metformin or a metformin containing product AND 2) Trial and failure, contraindication or intolerance to a medication from one of the following drug classes: sulfonylurea (e.g., glipizide, glimepiride), thiazolidinedione (e.g., pioglitazone), DPP-4 Inhibitor [e.g., Tradjenta (linagliptin)], SGLT2 inhibitor [e.g., Jardiance (empagliflozin)], GLP-1 receptor agonist [e.g., Trulicity (dulaglutide), Victoza (liraglutide)], or basal insulin (e.g., insulin glargine).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	DM (Reauth): Patient has experienced an objective response to therapy demonstrated by an improvement in HbA1c from baseline.

DAURISMO

Products Affected

- Daurismo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute myeloid leukemia (AML): Diagnosis of newly-diagnosed acute myeloid leukemia (AML) AND Daurismo therapy to be given in combination with low-dose cytarabine AND One of the following: 1) Patient is greater than or equal to 75 years old, or 2) Patient has significant comorbidities that preclude the use of intensive induction chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

DEFERASIROX

Products Affected

- Deferasirox

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Iron Overload due to Blood Transfusions (initial): Diagnosis of chronic iron overload (eg, sickle cell anemia, thalassemia, etc.) due to blood transfusion. Patient has blood transfusion of at least 100 mL/kg of packed red blood cells (eg, at least 20 units of packed red blood cells for a 40-kg person or more in individuals weighing more than 40 kg) prior to initiation of treatment with deferasirox. Patient has serum ferritin levels consistently greater than 1000 mcg/L prior to initiation of treatment with deferasirox. Chronic Overload in non-transfusion dependent thalassemia syndromes (initial): Diagnosis of chronic iron overload in non-transfusion dependent thalassemia syndrome. Patient has liver iron (Fe) concentration (LIC) levels consistently greater than or equal to 5 mg Fe per gram of dry weight prior to initiation of treatment with deferasirox. Patient has serum ferritin levels consistently more than 300 mcg/L prior to initiation of treatment with deferasirox. Chronic Iron Overload due to Blood Transfusions, Chronic Overload in non-transfusion dependent thalassemia syndromes (reauthorization): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by a hematologist/oncologist or hepatologist.
Coverage Duration	Plan year
Other Criteria	N/A

DEFERIPRONE

Products Affected

- Deferiprone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Transfusional iron overload: Diagnosis of transfusional iron overload due to one of the following: thalassemia syndromes, sickle cell disease, other transfusional-dependent anemia. Absolute Neutrophil Count (ANC) greater than $1.5 \times 10^9/L$. Trial and failure, contraindication or intolerance to one chelation therapy (e.g., generic deferasirox).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	All uses (reauth): Patient demonstrates positive clinical response to therapy. ANC greater than $1.5 \times 10^9/L$.

DEGARELIX

Products Affected

- Firmagon INJ 120MG/VIAL, 80MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

DICLOFENAC GEL 3%

Products Affected

- Diclofenac Sodium GEL 3%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Actinic Keratosis (initial): Diagnosis of Actinic Keratosis. Actinic Keratosis (reauthorization): Patient demonstrates positive clinical response to therapy. At least 30 days have elapsed since cessation of diclofenac sodium 3% topical gel therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	90 days
Other Criteria	N/A

DICLOFENAC TOPICAL SOLUTION

Products Affected

- Diclofenac Sodium EXTERNAL SOLN 1.5%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diclofenac topical solution 1.5%: Treatment of signs and symptoms of osteoarthritis of the knee(s).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

DIHYDROERGOTAMINE NASAL

Products Affected

- Dihydroergotamine Mesylate
NASAL SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of migraine headaches with or without aura. Will be used for the acute treatment of migraine. One of the following: Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or contraindication to all triptans.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy.

DOPTELET

Products Affected

- Doptelet

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Thrombocytopenia Prior to Planned Procedure (TPPP): Diagnosis (dx) of thrombocytopenia. Patient has chronic liver disease and is scheduled to undergo a procedure. Baseline platelet count is less than 50,000/mcL. Chronic Immune Thrombocytopenia (ITP) (initial): Diagnosis of chronic ITP or relapsed/refractory ITP. Baseline platelet count is less than 30,000/mcL. Trial and failure, contraindication, or intolerance to at least one of the following: corticosteroids (e.g., prednisone, methylprednisolone), immunoglobulins [e.g., Gammagard, immune globulin (human)], or splenectomy. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	TPPP: 1 month. ITP (initial, reauth): Plan year
Other Criteria	ITP (reauth): Patient demonstrates positive clinical response to therapy.

DOXEPIN TOPICAL

Products Affected

- Doxepin Hydrochloride CREA

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of moderate pruritus. Patient has atopic dermatitis or lichen simplex chronicus. Trial and failure, contraindication, or intolerance to at least one medium potency topical corticosteroid, or is not a candidate for topical corticosteroids (e.g., treatment is on face, axilla, or groin).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	14 days
Other Criteria	N/A

DRONABINOL

Products Affected

- Dronabinol

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Nausea and Vomiting Associated with Cancer Chemotherapy (CINV): Patient is receiving cancer chemotherapy. Trial and failure, contraindication, or intolerance (TF/C/I) to a 5HT-3 receptor antagonist (eg, Anzemet [dolasetron], Kytril [granisetron], or Zofran [ondansetron]). TF/C/I to one of the following: Ativan (lorazepam), Compazine (prochlorperazine), Decadron (dexamethasone), Haldol (haloperidol), Phenergan (promethazine), Reglan (metoclopramide), Zyprexa (olanzapine). AIDS anorexia: Diagnosis of anorexia with weight loss in patients with AIDS.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CINV: 6 months. AIDS anorexia: 3 months.
Other Criteria	Subject to Part B vs. Part D review. CINV: Approve for continuation of therapy for treatment covered under Part B when patient is receiving cancer chemotherapy.

DROXIDOPA

Products Affected

- Droxidopa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neurogenic orthostatic hypotension (NOH): (Initial): Diagnosis of symptomatic NOH. NOH is caused by one of the following conditions: primary autonomic failure (eg, Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, non-diabetic autonomic neuropathy. (Reauth): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	NOH (init): Prescribed by or in consultation with one of the following specialists: cardiologist, neurologist, nephrologist.
Coverage Duration	Initial: 1 month. Reauth: plan year
Other Criteria	Trial and failure, contraindication, or intolerance to one of the following agents: Fludrocortisone acetate, midodrine.

DUPIXENT

Products Affected

- Dupixent

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Atopic Dermatitis (AD) (initial): Diagnosis (dx) of moderate-to-severe AD. One of the following: a) Involvement of at least 10% body surface area (BSA), or b) SCORing Atopic Dermatitis (SCORAD) index value of at least 25. Trial and failure of a minimum 30-day supply (14-day supply for topical corticosteroids), contraindication (eg, safety concerns, not indicated for patient's age/weight), or intolerance to at least one of the following: a) Medium or higher potency topical corticosteroid, b) Pimecrolimus cream, c) Tacrolimus ointment, or d) Eucrisa (crisaborole) ointment. Eosinophilic Asthma (EA) (initial): Dx of moderate to severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-dupilumab treatment) peripheral blood eosinophil level greater than or equal to 150 cells/microliter. One of the following: 1) Patient has had two or more asthma exacerbations requiring systemic corticosteroids (eg, prednisone) within the past 12 mo, or 2) Prior asthma-related hospitalization within the past 12 mo. Corticosteroid Dependent Asthma (CDA) (initial): Dx of moderate to severe asthma. Patient is currently dependent on oral corticosteroids for the treatment of asthma. EA, CDA (initial): Patient is currently being treated with ONE of the following unless there is a contraindication or intolerance to these medications: 1) High-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and additional asthma controller medication [e.g., leukotriene receptor antagonist (eg, montelukast), long-acting beta-2 agonist (LABA) (eg, salmeterol), tiotropium] OR 2) One max-dosed combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Symbicort [budesonide/formoterol], Breo Ellipta (fluticasone/vilanterol)). Prurigo nodularis (PN) (init): Diagnosis of PN. Trial and failure, contraindication, or intolerance to one medium or higher potency topical corticosteroid.
Age Restrictions	Asthma (initial): Patient is 6 years of age or older. AD (initial): Patient is 6 months of age or older. CRSwNP, PN: no age restriction. EoE (initial): Patient is 1 year of age or older.
Prescriber Restrictions	AD, PN (init): Prescribed by or in consultation with a dermatologist or allergist/immunologist. Asthma (init): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. CRSwNP (initial): Prescribed by or in consultation with an otolaryngologist,

	allergist/immunologist, or pulmonologist. EoE (initial): Prescribed by or in consultation with a gastroenterologist or allergist/immunologist
Coverage Duration	CRSwNP, EoE (Init/Reauth): Plan year. Asthma, AD, PN (Init): 6 mo. Asthma, AD, PN (reauth): Plan yr.
Other Criteria	Eosinophilic esophagitis (EoE) (initial): Dx of EoE. Patient weighs at least 15 kg. Trial and failure, contraindication, or intolerance to one of the following: a) proton pump inhibitors (eg, pantoprazole, omeprazole) or b) topical (esophageal) corticosteroids (eg, budesonide, fluticasone). AD (reauth): Patient demonstrates positive clinical response to therapy. Chronic rhinosinusitis with nasal polyposis (CRSwNP) (initial): Diagnosis of CRSwNP. Unless contraindicated, the patient has had an inadequate response to 2 months of treatment with an intranasal corticosteroid (eg, fluticasone, mometasone). Used in combination with another agent for CRSwNP. EA (reauth): Patient demonstrates positive clinical response to therapy. CDA (reauth): Patient demonstrates positive clinical response to therapy. EA, CDA (reauth): Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications. CRSwNP (reauth): Patient demonstrates positive clinical response to therapy. Used in combination with another agent for CRSwNP. EoE, PN (reauth): Patient demonstrates positive clinical response to therapy.

ELIGARD

Products Affected

- Eligard

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Treatment of advanced prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

EMGALITY

Products Affected

- Emgality

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Episodic Migraines (EM) (120 mg/mL strength only) (initial): Diagnosis of EM. Patient has greater than or equal to 4 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) Topiramate (Topamax), e) Venlafaxine (Effexor), f) Candesartan (Atacand). Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines. Chronic Migraines (CM) (120 mg/mL strength only) (initial): Diagnosis of CM. Patient has greater than or equal to 8 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) OnabotulinumtoxinA (Botox), e) Topiramate (Topamax), f) Venlafaxine (Effexor), g) Candesartan (Atacand). Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines. Episodic Cluster Headaches (ECH) (100 mg/mL strength only) (initial): Diagnosis of ECH. Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months. Medication will not be used in combination with another injectable CGRP inhibitor.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	EM, CM, ECH (initial, reauth): Plan year.
Other Criteria	ECH (reauth): Patient demonstrates positive clinical response to therapy. Medication will not be used in combination with another injectable CGRP inhibitor. EM, CM (reauth): Patient demonstrates positive clinical

	response to therapy. Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.
--	--

ENBREL

Products Affected

- Enbrel INJ 25MG/0.5ML,
50MG/ML
- Enbrel Mini
- Enbrel Sureclick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of moderately to severely active PJIA. Minimum duration of a 6-week trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Plaque psoriasis (Initial): Diagnosis of moderate to severe chronic plaque psoriasis. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week TF/C/I to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Ankylosing Spondylitis (AS) (Initial): Diagnosis of active AS. Minimum duration of a one-month TF/C/I to one nonsteroidal anti-inflammatory drug (NSAID) (eg, ibuprofen, naproxen) at maximally tolerated doses.
Age Restrictions	N/A
Prescriber Restrictions	RA, JIA, AS (Initial): Prescribed by or in consultation with a rheumatologist. PsA (Initial): Prescribed by or in consultation with a rheumatologist or dermatologist. Plaque Psoriasis (Initial): Prescribed by or in consultation with a dermatologist.
Coverage Duration	All indications (initial): 6 months, (reauth): plan year
Other Criteria	All indications (Reauth): Patient demonstrates positive clinical response to therapy.

ENDARI

Products Affected

- L-glutamine PACK

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Sickle cell disease (initial): Diagnosis of sickle cell disease. Used to reduce acute complications of sickle cell disease. Patient has had 2 or more painful sickle cell crises within the past 12 months.
Age Restrictions	N/A
Prescriber Restrictions	Sickle cell disease (initial): Prescribed by or in consultation with a hematologist/oncologist
Coverage Duration	Sickle cell disease (initial, reauth): plan year
Other Criteria	Sickle cell disease (reauth): Patient demonstrates positive clinical response to therapy.

EPIDIOLEX

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Lennox-Gastaut syndrome (LGS): Diagnosis of seizures associated with LGS. Trial of, contraindication, or intolerance to two formulary anticonvulsants (e.g., topiramate, lamotrigine, valproate). Dravet syndrome (DS): Diagnosis of seizures associated with DS. Tuberous sclerosis complex (TSC): Diagnosis of seizures associated with TSC.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

EPOETIN ALFA (PREFERRED)

Products Affected

- Procrit
- Retacrit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Anemia due to Chronic Kidney Disease (CKD) (Initial): Diagnosis (dx) of CKD. Anemia by lab values (Hct less than 30% or Hgb less than 10 g/dL) within 30 days of request. The rate of hemoglobin decline indicates the likelihood of requiring a red blood cell (RBC) transfusion, and reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal. (Reauth): Diagnosis of CKD. Most recent or average (avg) Hct over 3 months is 33% or less (Hgb 11 g/dL or less) for patients on dialysis, without ESRD OR Most recent or average (avg) Hct over 3 months is 30% or less (Hgb 10 g/dL or less) for patients not on dialysis OR Most recent or average (avg) Hct over 3 months is 36% or less (Hgb 12 g/dL or less) for pediatric patients. Documentation of a positive clinical response to therapy from pre-treatment level. Anemia w/ HIV (Initial): Anemia by labs (Hgb less than 12 g/dL or Hct less than 36%) within 30 days of request. Serum erythropoietin level less than or equal to 500 mU/mL. Receiving zidovudine therapy or dx of HIV. (Reauth): Most recent or avg Hct over 3 months is below 36% or most recent or avg Hgb over 3 months is below 12 g/dl. Documentation of a positive clinical response to therapy from pre-treatment level. Anemia with chemo (Initial):Other causes of anemia ruled out. Anemia with labs (Hct less than 30%, Hgb less than 10 g/dL) within prior 2 weeks of request. Cancer is non-myeloid malignancy. Patient is receiving chemo. (Reauth): Anemia by labs (Hgb less than 10 g/dl or Hct less than 30%) collected within the prior 2 wks of request. Documentation of a positive clinical response to therapy from pre-treatment level. Patient is receiving chemo.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CKD,HIV(Init):6mo.(reauth):plan yr.Chemo(init, reauth):3mo.MDS(init):3mo,(reauth):plan yr.Preop:1 mo.
Other Criteria	Anemia in Myelodysplastic Syndrome (MDS) (Init): Diagnosis of MDS. Serum erythropoietin 500 mU/mL or less, or transfusion dependent MDS. (Reauth): Most recent or avg Hct over 3 months was 36% or less, OR most recent or avg Hgb over 3 months is 12 g/dl or less. Patient

demonstrates positive clinical response to therapy from pre-treatment level. Preoperative for reduction of allogeneic blood transfusion: Patient is scheduled to undergo elective, non-cardiac, non-vascular surgery. Hgb is greater than 10 to less than or equal to 13 g/dL. Patient is at high risk for perioperative transfusions. Patient is unwilling or unable to donate autologous blood pre-operatively. ESRD patients: Coverage is excluded under Medicare Part D for patients with ESRD on dialysis for any indication related or unrelated to treatment of ESRD since the payment for the drug is included in the ESRD PPS payment bundle. NON-ESRD PATIENTS for the following indications: Off-label uses (except Anemia in MDS and Hep C patients being treated with combo ribavirin and interferon/peginterferon): Will not be approved if patient has Hgb greater than 10 g/dL or Hct greater than 30%.CKD (init, reauth), Chemo (init), preop, MDS (init): Verify Fe eval for adequate Fe stores.

ERGOLOID

Products Affected

- Ergoloid Mesylates TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of one of the following: age-related dementia or Alzheimer's dementia. Trial and failure, contraindication, or intolerance to one of the following: a) donepezil, b) galantamine, or c) rivastigmine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Metastatic basal cell carcinoma (BCC): Diagnosis of metastatic basal cell carcinoma. Advanced basal cell carcinoma: Diagnosis of locally advanced basal cell carcinoma. One of the following: Cancer has recurred following surgery, Patient is not a candidate for surgery, or Patient is not a candidate for radiation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Non-metastatic castration-resistant prostate cancer (NM-CRPC): Diagnosis of non-metastatic, castration-resistant (chemical or surgical) or recurrent prostate cancer. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., Zoladex [goserelin], Vantas [histrelin], Lupron [leuprolide], Trelstar [triptorelin], Firmagon [degarelix]) OR 2) Patient received a bilateral orchiectomy.</p> <p>Metastatic castration-sensitive prostate cancer (M-CSPC): Diagnosis of metastatic, castration-sensitive or naive prostate cancer. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., Lupron[leuprolide], Trelstar[triptorelin], Zoladex [goserelin], Vantas [histrelin], Firmagon [degarelix]) OR 2) Patient received bilateral orchiectomy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

ERLOTINIB

Products Affected

- Erlotinib Hydrochloride TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): 1) Diagnosis of NSCLC. Disease is one of the following: a) advanced, b) metastatic, or c) recurrent. Tumors are positive for epidermal growth factor receptor (EGFR) exon 19 deletion mutations or exon 21 (L858R) substitution mutations or a known sensitizing EGFR mutation (e.g., in-frame exon 20 insertions, exon 18 G719 mutation, exon 21 L861Q mutation). Pancreatic cancer: Diagnosis of locally advanced, unresectable or metastatic pancreatic cancer. Used in combination with Gemzar (gemcitabine).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

EVEROLIMUS

Products Affected

- Everolimus TABS 10MG, 2.5MG, 5MG, 7.5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced Neuroendocrine Tumors (NET): Diagnosis (Dx) of neuroendocrine tumors of pancreatic origin, gastrointestinal origin, lung origin, or thymic origin. Disease is progressive. Disease is unresectable, locally advanced or metastatic. Advanced Renal Cell Carcinoma/Kidney Cancer: Dx of advanced renal cell cancer/kidney cancer. Disease is one of the following: (1) relapsed or (2) stage IV disease. Renal angiomyolipoma with tuberous sclerosis complex (TSC): Dx of renal angiomyolipoma and TSC, not requiring immediate surgery. Subependymal Giant Cell Astrocytoma (SEGA) with tuberous sclerosis (TS): Dx of SEGA associated with TS. Patient is not a candidate for curative surgical resection. Breast Cancer: Dx of recurrent or metastatic breast cancer. One of the following: Disease is hormone receptor positive (HR+) [i.e., estrogen-receptor-positive (ER+) or progesterone-receptor-positive (PR+)] OR both of the following: disease is HR- and disease has clinical characteristics that predict a HR+ tumor. Disease is HER2-negative. One of the following: Patient is a postmenopausal woman, patient is a premenopausal woman being treated with ovarian ablation/suppression, or patient is male. One of the following: A) Both of the following: a) one of the following: 1) Disease progressed while on or within 12 months of non-steroidal aromatase inhibitor [e.g., Arimidex (anastrozole), Femara (letrozole)] therapy or 2) patient was treated with tamoxifen at any time AND b) Used in combination with Aromasin (exemestane) OR B) Used in combination with Fulvestrant or Tamoxifen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	All uses: Approve for continuation of prior therapy if within the past 120 days.

EVEROLIMUS SOLUTION

Products Affected

- Everolimus TBSO

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced Neuroendocrine Tumors (NET): Diagnosis (Dx) of neuroendocrine tumors of pancreatic origin, gastrointestinal origin, lung origin, or thymic origin. Disease is progressive. Disease is unresectable, locally advanced or metastatic. Advanced Renal Cell Carcinoma/Kidney Cancer: Dx of advanced renal cell cancer/kidney cancer. Disease is one of the following: (1) relapsed or (2) stage IV disease. Renal angiomyolipoma with tuberous sclerosis complex (TSC): Dx of renal angiomyolipoma and TSC, not requiring immediate surgery. Subependymal Giant Cell Astrocytoma (SEGA) with tuberous sclerosis (TS): Dx of SEGA associated with TS. Patient is not a candidate for curative surgical resection. Breast Cancer: Dx of recurrent or metastatic breast cancer. One of the following: Disease is hormone receptor positive (HR+) [i.e., estrogen-receptor-positive (ER+) or progesterone-receptor-positive (PR+)] OR both of the following: disease is HR- and disease has clinical characteristics that predict a HR+ tumor. Disease is HER2-negative. One of the following: Patient is a postmenopausal woman, patient is a premenopausal woman being treated with ovarian ablation/suppression, or patient is male. One of the following: A) Both of the following: a) one of the following: 1) Disease progressed while on or within 12 months of non-steroidal aromatase inhibitor [e.g., Arimidex (anastrozole), Femara (letrozole)] therapy or 2) patient was treated with tamoxifen at any time AND b) Used in combination with Aromasin (exemestane) OR B) Used in combination with Fulvestrant or Tamoxifen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	TSC Associated Partial-Onset Seizures: Dx of TSC associated partial-onset seizures. Used as adjunctive therapy. All uses: Approve for continuation of prior therapy if within the past 120 days.

FASENRA

Products Affected

- Fasenra
- Fasenra Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of severe asthma. Asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter. One of the following: a) Patient has had two or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months, OR b) Prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications: a) Both of the following: 1) High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day) and 2) additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) OR b) One maximally-dosed combination ICS/LABA product [e.g., Advair (fluticasone propionate/salmeterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)].
Age Restrictions	Initial: Patient is 6 years of age or older
Prescriber Restrictions	Initial: Prescribed by or in consultation with a pulmonologist or allergy/immunology specialist
Coverage Duration	Initial: 6 mo, Reauth: Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications.

FENTANYL (PREFERRED)

Products Affected

- Fentanyl Citrate Oral Transmucosal

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cancer pain: Chart documentation provided reflecting oral transmucosal fentanyl will be used to manage pain related to an active cancer diagnosis. At least a one week history of one of the following medications to demonstrate tolerance to opioids: morphine sulfate at doses of greater than or equal to 60 mg/day, fentanyl transdermal patch at doses greater than or equal to 25 µg/hr, oxycodone at a dose of greater than or equal to 30 mg/day , oral hydromorphone at a dose of greater than or equal to 8 mg/day, oral oxymorphone at a dose of greater than or equal to 25 mg/day, an alternative opioid at an equianalgesic dose (eg, oral methadone greater than or equal to 20 mg/day). The patient is currently taking a long-acting opioid around the clock for cancer pain
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, pain specialist, hematologist, hospice care specialist, or palliative care specialist.
Coverage Duration	Plan year
Other Criteria	N/A

FINTEPLA

Products Affected

- Fintepla

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: 1) Diagnosis of seizures associated with Dravet syndrome, OR 2) Diagnosis of seizures associated with Lennox-Gastaut syndrome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

FLECTOR

Products Affected

- Diclofenac Epolamine

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Pain: Topical treatment of acute pain due to one of the following: minor strain, sprain, contusion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

FLUCYTOSINE

Products Affected

- Flucytosine CAPS 250MG, 500MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of one of the following: a) candidal endocarditis, b) candidiasis, c) candidiasis of urogenital site, d) candida endophthalmitis, e) central nervous system candidiasis, f) cryptococcosis, g) cryptococcal meningitis – HIV infection, h) HIV infection – pulmonary cryptococcosis, i) peritoneal dialysis-associated peritonitis, Fungal. Used in combination with amphotericin B.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 weeks
Other Criteria	N/A

FOTIVDA

Products Affected

- Fotivda

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of advanced renal cell carcinoma. Disease is one of the following: relapsed or refractory. Patient has received two or more prior systemic therapies (e.g., chemotherapy).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

FRUZAQLA

Products Affected

- Fruzaqla

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of metastatic colorectal cancer. Patient has been previously treated with both of the following: A) Fluoropyrimidine-, oxaliplatin-, irinotecan-based chemotherapy, and B) Anti-VEGF biological therapy (e.g., bevacizumab, ramucirumab). One of the following: A) Patient does not have RAS wild type tumors, OR B) Both of the following: a) Patient has RAS wild type tumors, AND b) Trial and failure, contraindication, or intolerance to both of the following: 1) An anti-EGFR biological therapy (e.g., panitumumab, cetuximab), and 2) One of the following: i) Trifluridine/tipiracil or ii) Regorafenib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

GAVRETO

Products Affected

- Gavreto

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is one of the following: recurrent, advanced, or metastatic. Presence of RET (rearranged during transfection) gene fusion-positive or RET rearrangement positive tumors. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is one of the following: advanced or metastatic. Disease is rearranged during transfection (RET) gene fusion-positive. Disease requires treatment with systemic therapy. One of the following: patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

GILOTRIF

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC. One of the following: 1) Tumors have non-resistant epidermal growth factor (EGFR) mutations as detected by an FDA-approved test OR 2) squamous disease progressing after previous platinum-based chemotherapy (e.g., cisplatin, carboplatin) OR 3) tumors are positive for a known sensitizing EGFR mutation (e.g., in-frame exon 20 insertions, exon 18 G719 mutation, exon 21 L861Q mutation).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

GLYCOPYRROLATE ORAL SOLUTION

Products Affected

- Glycopyrrolate ORAL SOLN

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic severe drooling: Diagnosis of chronic severe drooling (sialorrhea). Diagnosis of a neurologic condition (e.g., cerebral palsy, mental retardation, Parkinson disease) associated with chronic severe drooling (sialorrhea).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

GROWTH HORMONES

Products Affected

- Genotropin
- Genotropin Miniquick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>PGHD(initial):PGHD dx [confrmd by ht (utilizing age and gender grwth charts related to ht) documented(doc) by ht more than 2.0SD below midparental ht or more than 2.25SD below population(pop) mean (below 1.2 percentile for age and gender),or grwth velocity more than 2SD below mean for age and gender, or delayed skeletal maturation more than 2SD below mean for age and gender (eg,delayed more than 2yrs compared w/chronological age)]. PWS(reauth):evidence of positive response to tx(eg,incr in total LBM, decr in fat mass) and expctd adult ht not attained and doc of expctd adult ht goal. GFSGA(initial):SGA dx based on catchup grwth failure in 1st 24mo of life using 0-36mo grwth chart confrmd by birth wt or length below 3rd percentile for gestational age(more than 2SD below pop mean) and ht remains at or below 3rd percentile (more than 2SD below pop mean). TS,NS(initial):ped grwth failure dx assoc w/TS w/doc female w/bone age less than 14yrs, or NS and ht below 5th percentile on grwth charts for age and gender. SHOX(initial):ped grwth failure dx w/SHOX gene deficiency confirmed by genetic testing. GFCRI(initial): ped grwth failure dx assoc w/CRI. ISS(initial):ISS dx, diagnostic eval excluded other causes assoc w/short stature(eg GHD, chronic renal insufficiency), doc ht at or below -2.25SD score below corresponding mean ht for age and gender assoc with growth rates unlikely to permit attainment of adult height in the normal range. PGHD,NS,SHOX,GFCRI,ISS (initial): doc male w/bone age less than 16yrs or female w/bone age less than 14yrs. PGHD,GFSGA,TS/NS,SHOX,GFCRI,ISS(reauth):expctd adult ht not attained and doc of expctd adult ht goal.</p>
Age Restrictions	N/A
Prescriber Restrictions	PGHD, PWS, GFSGA, TS/NS, SHOX, AGHD, IGHDA, ISS: prescribed by or in consultation with an endocrinologist. GFCRI: prescribed by or in consultation with an endocrinologist or nephrologist
Coverage Duration	All indications (initial, reauth): Plan year
Other Criteria	AGHD(initial):dx of AGHD as a result of clin records supportng dx of childhood-onset GHD, or adult-onset GHD w/clin records doc hormone

deficiency d/t hypothalamic-pituitary dz from organic or known causes (eg,damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and pt has 1GH stim test (insulin tolerance test [ITT],glucagon,macimorelin) to confirm adult GHD w/corresponding peak GH values ([ITT at or below 5mcg/L],[glucagon at or below 3mcg/L],[macimorelin less than 2.8 ng/mL 30, 45, 60 and 90 mins after admin]) or doc deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IGF-1/somatomedinC below age and gender adjstd nrml range as provided by physicians lab.
AGHD,IGHDA(reauth):evidence of ongoing monitoring as demonstrated by doc w/in past 12mo of IGF-1/somatomedinC level.
IGHDA(initial):doc GHD after 2 GH stim tests(ITT,glucagon,macimorelin) w/ 2 corresponding peak GH values [ITT at or below 5mcg/L],[glucagon at or below 3mcg/L],[macimorelin less than 2.8 ng/mL 30,45,60,90 mins after admin].

HAEGARDA

Products Affected

- Haegarda

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prophylaxis of hereditary angioedema (HAE) attacks (initial): Diagnosis of HAE. Diagnosis has been confirmed by C1 inhibitor (C1-INh) deficiency or dysfunction (Type I or II HAE) as documented by ONE of the following: a) C1-INH antigenic level below the lower limit of normal OR b) C1-INH functional level below the lower limit of normal. For prophylaxis against HAE attacks. Not used in combination with other approved treatments for prophylaxis against HAE attacks.
Age Restrictions	N/A
Prescriber Restrictions	HAE (prophylaxis) (initial): Prescribed by or in consultation with an immunologist or an allergist
Coverage Duration	HAE (prophylaxis) (initial, reauth): Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. Not used in combination with other approved treatments for prophylaxis against HAE attacks

HUMIRA

Products Affected

- Humira INJ 10MG/0.1ML, 20MG/0.2ML, 40MG/0.4ML, 40MG/0.8ML
- Humira Pen
- Humira Pen-cd/uc/hs Starter INJ 80MG/0.8ML
- Humira Pen-pediatric Uc Starter Pack
- Humira Pen-ps/uv Starter INJ 0

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Rheumatoid Arthritis (RA) (Initial): Diagnosis (Dx) of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate (MTX), leflunomide, sulfasalazine. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Dx of moderately to severely active PJIA. Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. Psoriatic Arthritis (PsA) (Initial): Dx of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Plaque psoriasis (PsO) (Initial): Dx of moderate to severe chronic PSO. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week TF/C/I to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Ankylosing Spondylitis (AS) (Initial): Dx of active AS. Minimum duration of a one-month TF/C/I to one NSAID (eg, ibuprofen, naproxen) at maximally tolerated doses. Crohn's Disease (CD) (Initial): Dx of moderately to severely active CD. One of the following: frequent diarrhea and abdominal pain, at least 10% weight loss, complications (eg, obstruction, fever, abdominal mass), abnormal lab values (eg, CRP), OR CD Activity Index (CDAI) greater than 220. TF/C/I to one of the following conventional therapies: 6-mercaptopurine (6-MP), azathioprine, corticosteroid (eg, prednisone), MTX. Uveitis (UV) (Initial): Dx of non-infectious uveitis classified as intermediate, posterior, or panuveitis.</p>
Age Restrictions	N/A
Prescriber Restrictions	RA, AS, JIA: (Initial) Prescribed by or in consultation with a rheumatologist. PsA: (Initial) Prescribed by or in consultation with a dermatologist or rheumatologist. Plaque Psoriasis, HS: (Initial) Prescribed by or in consultation with a dermatologist. CD, UC: (Initial) Prescribed by or in consultation with a gastroenterologist. UV (initial): Prescribed by or in consultation with a rheumatologist or an ophthalmologist.

Coverage Duration	UC: (Initial) 12 wks. Other uses (initial): 6 months. All uses (reauth): plan year.
Other Criteria	<p>Ulcerative Colitis (UC) (Initial): Dx of moderately to severely active UC. One of the following: greater than 6 stools per day, frequent blood in the stools, frequent urgency, presence of ulcers, abnormal lab values (eg, hemoglobin, ESR, CRP), OR dependent on, or refractory to, corticosteroids. TF/C/I to one of the following conventional therapies: 6-MP, azathioprine, corticosteroid (eg, prednisone), aminosalicylate (eg, mesalamine, olsalazine, sulfasalazine). Hidradenitis suppurativa (HS) (Initial): Dx of moderate to severe hidradenitis suppurativa (ie, Hurley Stage II or III). RA, PJA, PsA, AS, PsO, CD, HS, UV (Reauth): Patient demonstrates positive clinical response to therapy. UC (Reauth): For patients who initiated therapy within the past 12 weeks: Patient has clinical remission or significant clinical benefit by eight weeks (Day 57) of therapy OR For patients who have been maintained on therapy for longer than 12 weeks: Patient demonstrates positive clinical response to therapy.</p>

IBRANCE

Products Affected

- Ibrance

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of breast cancer. Disease is a) locally advanced, metastatic, recurrent, or Stage IV
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ICATIBANT

Products Affected

- Icatibant Acetate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Treatment of hereditary angioedema (HAE) attacks (initial): Diagnosis of HAE. Diagnosis has been confirmed by C1 inhibitor (C1-INh) deficiency or dysfunction (Type I or II HAE) as documented by ONE of the following: a) C1-INH antigenic level below the lower limit of normal OR b) C1-INH functional level below the lower limit of normal. For the treatment of acute HAE attacks. Not used in combination with other approved treatments for acute HAE attacks.
Age Restrictions	N/A
Prescriber Restrictions	HAE (initial): Prescribed by or in consultation with an immunologist or an allergist
Coverage Duration	Initial, Reauth: Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. Not used in combination with other approved treatments for acute HAE attacks

ICLUSIG

Products Affected

- Iclusig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Myelogenous / Myeloid Leukemia (CML): Diagnosis of chronic myelogenous/myeloid leukemia (CML). One of the following: a) Both of the following: Disease is in the chronic phase AND patient is unable to take or has failed treatment with two or more alternative tyrosine kinase inhibitors (TKI) [eg, Bosulif (bosutinib), imatinib, Sprycel (dasatinib), Tasigna (nilotinib)], b) confirmed documentation of T315I mutation, or c) Both of the following: Disease is in the accelerated or blast phase AND no other kinase inhibitors are indicated. Acute Lymphoblastic Leukemia: Diagnosis of Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ALL). One of the following: a) No other kinase inhibitors are indicated, b) Confirmed documentation of T315I mutation, c) Disease is relapsed or refractory, d) Used as a component of HyperCVAD regimen (hyper-fractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone, alternating with high-dose methotrexate and cytarabine) induction or consolidation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

IDHIFA

Products Affected

- Idhifa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of AML. AML is isocitrate dehydrogenase-2 (IDH2) mutation-positive. One of the following: A) Disease is relapsed or refractory, B) Patient is not a candidate for or patient declines intensive induction therapy, C) Used for post induction therapy following response to low intensity induction therapy, or D) Used for consolidation therapy as continuation of low-intensity regimen used for induction.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

IMATINIB

Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Myelogenous/Myeloid Leukemia (CML): Diagnosis of CML (CML). Acute Lymphoblastic Leukemia (ALL): Diagnosis of Philadelphia chromosome positive/BCR ABL-positive ALL (Ph+/BCR ABL+ ALL). Myelodysplastic/ myeloproliferative disease (MDS/MPD): Diagnosis of MDS/MPD. One of the following: 1) Disease is associated with 5q32 translocations or 2) Disease is associated with platelet-derived growth factor receptor (PDGFR) gene rearrangements or 3) disease is associated with a t(5:12) translocation associated with the ETV6-PDGFRbeta fusion gene. Aggressive systemic mastocytosis (ASM): Diagnosis of ASM. Patient is without the D816V c-Kit mutation or c-Kit mutational status unknown or eosinophilia is present with FIP1L1-PDGFRalpha fusion gene. Hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL): Diagnosis of at least one of the following: HES or CEL. Dermatofibrosarcoma protuberans (DFSP): Diagnosis of DFSP. Gastrointestinal Stromal Tumors (GIST): Diagnosis of GIST.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

IMBRUVICA

Products Affected

- Imbruvica CAPS
- Imbruvica SUSP
- Imbruvica TABS 140MG, 280MG, 420MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Waldenstrom's Macroglobulinemia (WM)/Lymphoplasmacytic Lymphoma (LPL): Diagnosis of WM/LPL. Chronic Graft Versus Host Disease (cGVHD): Diagnosis of cGVHD AND trial and failure of at least one other systemic therapy (e.g., corticosteroids like prednisone or methylprednisolone, mycophenolate, etc.).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

IMVEXXY

Products Affected

- Imvexxy Maintenance Pack
- Imvexxy Starter Pack

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Dyspareunia: Diagnosis of moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

INBRIJA

Products Affected

- Inbrija

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Parkinson's disease (PD) (initial): Diagnosis of PD. Patient is experiencing intermittent OFF episodes. Used in combination with carbidopa/levodopa.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	PD (initial, reauth): Plan year
Other Criteria	PD (reauth): Patient demonstrates positive clinical response to therapy. Used in combination with carbidopa/levodopa.

INGREZZA

Products Affected

- Ingrezza

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Tardive Dyskinesia: Diagnosis of tardive dyskinesia. One of the following: a) patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication OR b) patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Chorea associated with Huntington's disease: Diagnosis of chorea in patients with Huntington's disease.
Age Restrictions	N/A
Prescriber Restrictions	Tardive Dyskinesia: Prescribed by or in consultation with a neurologist or psychiatrist. Chorea associated with Huntington's disease: Prescribed by or in consultation with a neurologist.
Coverage Duration	All Indications: Plan year
Other Criteria	N/A

INLYTA

Products Affected

- Inlyta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced Renal Cell Carcinoma: Diagnosis of renal cell cancer. One of the following: (1) diagnosis of stage IV disease, or (2) Both of the following: a) Disease is advanced, and b) One of the following: i) Patient has failed one prior systemic therapy (e.g., chemotherapy), or ii) Inlyta will be used in combination with Bavencio (avelumab) or Keytruda (pembrolizumab).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

INQOVI

Products Affected

- Inqovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: 1) Both of the following: a) Diagnosis of myelodysplastic syndrome (MDS) and b) Patient is intermediate-1, intermediate-2, or high-risk per the International Prognostic Scoring System (IPSS), OR 2) Diagnosis of chronic myelomonocytic leukemia (CMML).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

INREBIC

Products Affected

- Inrebic

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Myelofibrosis: Diagnosis of one of the following: primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

INSULIN - LIKE GROWTH FACTOR

Products Affected

- Increlex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial therapy: IGF-1 deficiency: Diagnosis of severe primary IGF-1 deficiency. Height standard deviation score of -3.0 or less. Basal IGF-1 standard deviation score of -3.0 or less. Normal or elevated growth hormone (GH). GH gene deletion: Diagnosis of growth hormone gene deletion who have developed neutralizing antibodies to GH. Reauthorization: Patient demonstrates positive clinical response to therapy. Both of the following: (1) Expected adult height is not obtained and (2) Documentation of expected adult height goal
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with an endocrinologist
Coverage Duration	Initial, reauth: plan year
Other Criteria	N/A

IRESSA

Products Affected

- Gefitinib

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic NSCLC, and One of the following: tumors are positive for epidermal growth factor receptor (EGFR) exon 19 deletions or tumors are positive for EGFR exon 21 (L858R) substitution mutations or tumors are positive for a known sensitizing EGFR mutation (e.g., in-frame exon 20 insertions, exon 18 G719 mutation, exon 21 L861Q mutation).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ISOTRETINOIN

Products Affected

- Accutane CAPS 10MG, 20MG, 40MG
- Amnesteem
- Claravis
- Isotretinoin CAPS
- Zenatane

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	(initial): Diagnosis of severe recalcitrant nodular acne unresponsive to conventional therapy OR diagnosis of treatment resistant acne. Trial and failure, contraindication or intolerance to an adequate trial (at least 6 weeks) on two of the following conventional therapy regimens: a) topical retinoid or retinoid-like agent [eg, Retin-A/Retin-A Micro (tretinoin)], b) oral antibiotic [eg, Ery-Tab (erythromycin), Minocin (minocycline)], c) topical antibiotic with or without benzoyl peroxide [eg, Cleocin-T (clindamycin), erythromycin, BenzaClin (benzoyl peroxide/clindamycin), Benzamycin (benzoyl peroxide/erythromycin)]. Retreatment (Reauthorization): After greater than or equal to 2 months off therapy, persistent or recurring severe recalcitrant nodular acne is still present.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 6 months. Reauth: Retreatment - 6 months
Other Criteria	N/A

ISTURISA

Products Affected

- Isturisa TABS 1MG, 5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cushing's disease (initial): Diagnosis of Cushing's disease. One of the following: a) Patient is not a candidate for pituitary surgery, OR b) Pituitary surgery has not been curative for the patient.
Age Restrictions	N/A
Prescriber Restrictions	Cushing's disease (initial): Prescribed by or in consultation with an endocrinologist.
Coverage Duration	Cushing's disease (initial, reauth): Plan Year
Other Criteria	Cushing's disease (reauth): Patient demonstrates positive clinical response to therapy.

ITRACONAZOLE (CAPSULES)

Products Affected

- Itraconazole CAPS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Systemic Fungal Infections: Diagnosis of blastomycosis, histoplasmosis, or aspergillosis. Onychomycosis: Diagnosis of fingernail or toenail onychomycosis confirmed by one of the following: KOH test, fungal culture, or nail biopsy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Systemic Fungal infxns: plan year. Onychomycosis: (Fingernail) 2 mo. (Toenail) 3 mo.
Other Criteria	N/A

IVERMECTIN TABLETS

Products Affected

- Ivermectin TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Any medically accepted indication [e.g., Onchocerciasis due to nematode parasite, Pediculosis, Strongyloidiasis, Ascariasis, Scabies (including crusted scabies), Cutaneous larva migrans (hook worm disease), Enterobiasis, Filariasis, Trichuriasis, or Gnathostomiasis].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

IVIG (NON - PREFERRED)

Products Affected

- Panzyga

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Primary immunodeficiency syndrome (PIS): patients with PIS. Idiopathic Thrombocytopenic Purpura (ITP): diagnosis (dx) of ITP. Documented (doc) platelet count of less than $50 \times 10^9/L$. Kawasaki disease (KD): dx of KD. B-cell Chronic Lymphocytic Leukemia (CLL): dx of B-Cell CLL. Doc hypogammaglobulinemia (IgG less than 500mg/dL) or history of bacterial infections associated with B-cell CLL. Bone Marrow Transplant (BMT): Confirmed allogeneic BMT within the last 100 days. Doc severe hypogammaglobulinemia (IgG less than 400 mg/dL). HIV:dx of HIV. 13 years of age or less. Doc hypogammaglobulinemia (IgG less than 400 mg/dL) or functional antibody deficiency demonstrated by poor specific antibody titers or recurrent bacterial infections. Guillain-Barre Syndrome (GBS) initial: dx of GBS. severe disease requiring aid to walk. Onset of neuropathic symptoms in the last 4 weeks. Myasthenia Gravis (MG): dx of generalized MG. Evidence of myasthenic exacerbation, defined by 1 of the following sx's in the last month: difficulty swallowing, acute respiratory failure, or major functional disability responsible for the discontinuation of physical activity. Concomitant immunomodulator therapy (tx)(eg, azathioprine, cyclosporine), unless contraindicated, will be used for long-term management of MG. Dermatomyositis and Polymyositis (D/P) initial: dx of dermatomyositis or polymyositis. Trial and failure, contraindication or intolerance (TF/C/I) to immunosuppressive tx (eg corticosteroids, methotrexate, azathioprine, cyclophosphamide). Stiff person syndrome (SPS) initial:dx of SPS. TF/C/I to GABAergic medication (eg, baclofen). TF/C/I to immunosuppressive tx (eg, azathioprine, corticosteroids). Lambert-Eaton myasthenic syndrome (LEMS) initial: dx of LEMS. TF/C/I to immunomodulator monotherapy (eg, azathioprine, corticosteroids). Concomitant immunomodulator tx (eg, azathioprine, corticosteroids), unless contraindicated, will be used for long-term management of LEMS.</p>
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g., immunologist, hematologist, neurologist).

Coverage Duration	KD: 1 mo. GBS,CIDP (initial), MG: 3 mo. ITP: 6 mo. CIDP,GBS (reauth), other uses: plan year.
Other Criteria	<p>All (except reauth): TF/C/I to one of the following: Bivigam, Flebogamma, Gammagard, Gammagard SD, Gammaked, Gammaplex, Gamunex-C, Octagam, Privigen, OR allow for continuation of therapy if within the past 120 days. Subject to Part B vs. Part D review. Pt does not meet criteria for Part B or patient is in a long-term care facility. PIS: Clinically significant functional deficiency of humoral immunity as evidenced by doc failure to produce antibodies to specific antigens or hx of significant recurrent infxns. Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) initial: dx of CIDP confirmed by: 1) progressive sxns present for at least 2 mo, 2) symptomatic polyradiculoneuropathy as indicated by progressive or relapsing motor impairment of more than 1 limb, OR Progressive or relapsing sensory impairment of more than 1 limb, 3) Electrophysiologic findings when 3 of the following 4 criteria are present: Partial conduction block of 1 or more motor nerve, Reduced conduction velocity of 2 or more motor nerves, Prolonged distal latency of 2 or more motor nerves, Prolonged F-wave latencies of 2 or more motor nerves or the absence of F waves. Multifocal motor neuropathy (MMN) initial: dx of MMN confirmed by all of the following: 1) weakness with slowly progressive or stepwise progressive course over at least 1 month, 2) asymmetric involvement of 2 or more nerves, AND 3) absence of motor neuron signs and bulbar signs. CIDP, MMN reauth: documentation of positive clinical response to tx as measured by an objective scale [eg, Rankin, Modified Rankin, Medical Research Council (MRC) scale]. Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect. Relapsing remitting Multiple Sclerosis (MS) initial: dx of relapsing remitting form of MS (RRMS). Documentation of an MS exacerbation or progression (worsening) of the patient's clinical status from the visit prior to the one prompting the decision to initiate immune globulin tx. TF/c/I to 2 of the following: teriflunomide, interferon beta-1b, interferon beta-1a, glatiramer acetate, natalizumab, dimethyl fumarate, fingolimod. RRMS reauth: The prescriber maintains and provides chart documentation of the patient's evaluation, including all of the following: findings of interval examination including neurological deficits incurred, and assessment of disability (eg, Expanded Disability Status Score [EDSS], Functional Systems Score [FSS], Multiple Sclerosis Functional Composite [MSFC], Disease Steps [DS]). Stable or improved disability score (eg, EDSS, FSS, MSFC, DS). Documentation of decreased number of relapses since starting immune globulin tx. Dx continues to be the relapsing-remitting form of MS. Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect. GBS, D/P, SPS, LEMS reauth:</p>

	Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect.
--	--

IVIG (PREFERRED)

Products Affected

- Bivigam INJ 5GM/50ML
- Gammagard Liquid INJ
2.5GM/25ML
- Gammagard S/d Iga Less Than
1mcg/ml
- Gammaked INJ 1GM/10ML
- Gammaplex INJ 10GM/100ML,
10GM/200ML, 20GM/200ML,
5GM/50ML
- Gamunex-c INJ 1GM/10ML
- Octagam INJ 1GM/20ML,
2GM/20ML
- Privigen INJ 20GM/200ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Primary immunodeficiency syndrome (PIS): patients with PIS. Idiopathic Thrombocytopenic Purpura (ITP): diagnosis (dx) of ITP. Documented (doc) platelet count of less than $50 \times 10^9/L$. Kawasaki disease (KD): dx of KD. B-cell Chronic Lymphocytic Leukemia (CLL): dx of B-Cell CLL. Doc hypogammaglobulinemia (IgG less than 500mg/dL) or history of bacterial infections associated with B-cell CLL. Bone Marrow Transplant (BMT): Confirmed allogeneic BMT within the last 100 days. Doc severe hypogammaglobulinemia (IgG less than 400 mg/dL). HIV:dx of HIV. 13 years of age or less. Doc hypogammaglobulinemia (IgG less than 400 mg/dL) or functional antibody deficiency demonstrated by poor specific antibody titers or recurrent bacterial infections. Guillain-Barre Syndrome (GBS) initial: dx of GBS. severe disease requiring aid to walk. Onset of neuropathic symptoms in the last 4 weeks. Myasthenia Gravis (MG): dx of generalized MG. Evidence of myasthenic exacerbation, defined by 1 of the following sx's in the last month: difficulty swallowing, acute respiratory failure, or major functional disability responsible for the discontinuation of physical activity. Concomitant immunomodulator therapy (tx)(eg, azathioprine, cyclosporine), unless contraindicated, will be used for long-term management of MG. Dermatomyositis and Polymyositis (D/P) initial: dx of dermatomyositis or polymyositis. Trial and failure, contraindication or intolerance (TF/C/I) to immunosuppressive tx (eg corticosteroids, methotrexate, azathioprine, cyclophosphamide). Stiff person syndrome (SPS) initial:dx of SPS. TF/C/I to GABAergic medication (eg, baclofen). TF/C/I to immunosuppressive tx (eg, azathioprine, corticosteroids). Lambert-Eaton myasthenic syndrome (LEMS) initial: dx of LEMS. TF/C/I to immunomodulator monotherapy (eg, azathioprine, corticosteroids). Concomitant immunomodulator tx (eg, azathioprine, corticosteroids), unless contraindicated, will be used for long-term management of LEMS.</p>
Age Restrictions	N/A
Prescriber Restrictions	All uses: Prescribed by or in consultation with a physician who has specialized expertise in managing patients on immune globulin therapy (e.g., immunologist, hematologist, neurologist).

Coverage Duration	KD: 1 mo. GBS,CIDP (initial), MG: 3 mo. ITP: 6 mo. CIDP,GBS (reauth), other uses: plan year.
Other Criteria	<p>Subject to Part B vs. Part D review. Pt does not meet criteria for Part B or patient is in a long-term care facility. PIS: Clinically significant functional deficiency of humoral immunity as evidenced by doc failure to produce antibodies to specific antigens or hx of significant recurrent infxns.</p> <p>Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) initial: dx of CIDP confirmed by: 1) progressive sxs present for at least 2 mo, 2) symptomatic polyradiculoneuropathy as indicated by progressive or relapsing motor impairment of more than 1 limb, OR Progressive or relapsing sensory impairment of more than 1 limb, 3) Electrophysiologic findings when 3 of the following 4 criteria are present: Partial conduction block of 1 or more motor nerve, Reduced conduction velocity of 2 or more motor nerves, Prolonged distal latency of 2 or more motor nerves, Prolonged F-wave latencies of 2 or more motor nerves or the absence of F waves. Multifocal motor neuropathy (MMN) initial: dx of MMN confirmed by all of the following: 1) weakness with slowly progressive or stepwise progressive course over at least 1 month, 2) asymmetric involvement of 2 or more nerves, AND 3) absence of motor neuron signs and bulbar signs. CIDP, MMN reauth: documentation of positive clinical response to tx as measured by an objective scale [eg, Rankin, Modified Rankin, Medical Research Council (MRC) scale]. Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect. Relapsing remitting Multiple Sclerosis (MS) initial: dx of relapsing remitting form of MS (RRMS). Documentation of an MS exacerbation or progression (worsening) of the patient's clinical status from the visit prior to the one prompting the decision to initiate immune globulin tx. TF/c/I to 2 of the following: teriflunomide, interferon beta-1b, interferon beta-1a, glatiramer acetate, natalizumab, dimethyl fumarate, fingolimod. RRMS reauth: The prescriber maintains and provides chart documentation of the patient's evaluation, including all of the following: findings of interval examination including neurological deficits incurred, and assessment of disability (eg, Expanded Disability Status Score [EDSS], Functional Systems Score [FSS], Multiple Sclerosis Functional Composite [MSFC], Disease Steps [DS]). Stable or improved disability score (eg, EDSS, FSS, MSFC, DS). Documentation of decreased number of relapses since starting immune globulin tx. Dx continues to be the relapsing-remitting form of MS. Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect. GBS, D/P, SPS, LEMS reauth: Documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect.</p>

IWILFIN

Products Affected

- Iwilfin

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of high-risk neuroblastoma (HRNB). Patient has shown at least a partial response to prior multiagent, multimodality therapy. Prior therapy included anti-GD2 immunotherapy (e.g., Danyelza [naxitamab-gqgk], Unituxin [dinutuximab]).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

JAKAFI

Products Affected

- Jakafi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Myelofibrosis: One of the following: Primary myelofibrosis, post-polycythemia vera myelofibrosis, or post-essential thrombocythemia myelofibrosis. Polycythemia vera: Diagnosis of polycythemia vera and trial and failure, contraindication, or intolerance to one of the following: hydroxyurea or interferon therapy (e.g., Intron A, pegasys, etc.). Graft versus host disease (GVHD): One of the following: A) Both of the following: Diagnosis of acute (aGVHD). Disease is steroid-refractory, or B) Both of the following: Diagnosis of chronic graft versus host disease (cGVHD). Trial and failure of at least one or more lines of systemic therapy (e.g., corticosteroids, mycophenolate).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

JAYPIRCA

Products Affected

- Jaypirca

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of mantle cell lymphoma (MCL). Disease is one of the following: a) relapsed, or b) refractory. Patient has received at least two prior systemic therapies [e.g., chemotherapy] for MCL, one of which is a Bruton Tyrosine Kinase (BTK) inhibitor therapy [e.g., Calquence (acalabrutinib), Imbruvica (ibrutinib)]. Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of one of the following: a) CLL, or b) SLL. Patient has received treatment for CLL/SLL with both of the following therapies: a) BTK inhibitor therapy [e.g., Calquence (acalabrutinib), Imbruvica (ibrutinib)], and b) B-cell lymphoma 2 (BCL-2) inhibitor therapy [e.g., Venclexta (venetoclax)].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

JYLAMVO

Products Affected

- Jylamvo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neoplastic diseases: Diagnosis of one of the following: A) acute lymphoblastic leukemia (ALL), B) mycosis fungoides (cutaneous T-cell lymphoma), or C) relapsed or refractory non-hodgkin lymphomas. Rheumatoid arthritis (RA) (initial): Diagnosis of RA. Psoriasis (initial): Diagnosis of severe psoriasis.
Age Restrictions	N/A
Prescriber Restrictions	RA: Prescribed by or in consultation with a rheumatologist. Psoriasis: Prescribed by or in consultation with a dermatologist.
Coverage Duration	Neoplastic diseases: Plan Year. RA, Psoriasis (initial, reauth): Plan Year.
Other Criteria	Neoplastic diseases: Approve for continuation of prior therapy within the past 120 days. RA, Psoriasis (reauth): Patient demonstrates positive clinical response to therapy.

KALYDECO

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic Fibrosis (CF) (Initial): Diagnosis of CF. Submission of laboratory records confirming patient has at least one mutation in the CFTR gene that is responsive to ivacaftor potentiation.
Age Restrictions	CF (initial): Patient is 1 month of age or older.
Prescriber Restrictions	CF (initial, reauth): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
Coverage Duration	CF (initial): 6 mos, (reauth): plan year
Other Criteria	CF (reauth): Patient demonstrates positive clinical response to therapy.

KERENDIA (SGLT2)

Products Affected

- Kerendia

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of chronic kidney disease (CKD) associated with type 2 diabetes (T2D). Urine albumin-to-creatinine ratio (UACR) greater than or equal to 30 mg/g. Estimated glomerular filtration rate (eGFR) greater than or equal to 25 mL/min/1.73 m ² . Serum potassium level less than or equal to 5.0 mEq/L prior to initiating treatment. One of the following: 1) Minimum 30-day supply trial of a maximally tolerated dose and will continue therapy with one of the following: a) generic angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril), or b) generic angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan), OR 2) Patient has a contraindication or intolerance to ACE inhibitors and ARBs. One of the following: 1) Patient is on a stable dose and will continue therapy with a sodium-glucose cotransporter-2 (SGLT2) inhibitor (e.g., Farxiga [dapagliflozin], Jardiance [empagliflozin]), OR 2) Trial and failure, contraindication or intolerance to an SGLT2 inhibitor (e.g., Farxiga [dapagliflozin], Jardiance [empagliflozin]).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, Reauth: Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. One of the following: 1) Patient continues to be on a maximally tolerated dose of ACE inhibitor or ARB, OR 2) Patient has a contraindication or intolerance to ACE inhibitors and ARBs.

KISQALI

Products Affected

- Kisqali

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of advanced, recurrent, or metastatic breast cancer. Cancer is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: A) Used in combination with an aromatase inhibitor (e.g., anastrozole, letrozole, exemestane) or B) Used in combination with Faslodex (fulvestrant).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

KISQALI - FEMARA PACK

Products Affected

- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of advanced, recurrent, or metastatic breast cancer. Cancer is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

KORLYM

Products Affected

- Mifepristone TABS 300MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	(Initial): Diagnosis of endogenous Cushing's syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Diagnosis of either type 2 diabetes mellitus or diagnosis of glucose intolerance. Patient has either failed surgery or patient is not a candidate for surgery. Patient is not pregnant. (Reauthorization): Documentation of one of the following: patient has improved glucose tolerance while on therapy or patient has stable glucose tolerance while on therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

KOSELUGO

Products Affected

- Koselugo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neurofibromatosis Type 1 (NF1): Diagnosis of NF1. Patient has plexiform neurofibromas that are both of the following: inoperable and causing significant morbidity (e.g., disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment, bladder/bowel dysfunction).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

KRAZATI

Products Affected

- Krazati

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): Diagnosis of NSCLC. Presence of KRAS G12C mutation. Disease is recurrent, advanced, or metastatic. Patient has received at least one prior systemic therapy (e.g., chemotherapy).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

LAPATINIB

Products Affected

- Lapatinib Ditosylate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer: One of the following: A) Diagnosis of recurrent or stage IV hormone receptor positive (HR+), human epidermal growth factor receptor 2-positive (HER2+) breast cancer. Used in combination with an aromatase inhibitor [eg, Aromasin (exemestane), Femara (letrozole), Arimidex (anastrozole)]. OR B) Diagnosis of recurrent or metastatic HER2+ breast cancer. Used in combination with trastuzumab or Xeloda (capecitabine).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LENVIMA

Products Affected

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Differentiated thyroid cancer (DTC): Diagnosis of DTC. Renal cell carcinoma: Diagnosis of Advanced Renal cell carcinoma. Hepatocellular Carcinoma (HCC): Diagnosis of Hepatocellular Carcinoma or liver cell carcinoma. Endometrial Carcinoma: Diagnosis of advanced endometrial carcinoma.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LEUPROLIDE ACETATE

Products Affected

- Leuprolide Acetate INJ 1MG/0.2ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Treatment of advanced prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prostate Cancer: plan year
Other Criteria	Prostate Cancer: Approve for continuation of prior therapy if within the past 120 days.

LIBERVANT

Products Affected

- Libervant

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Epilepsy: Diagnosis of epilepsy. Frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LIDOCAINE PATCH

Products Affected

- Lidocaine PTCH 5%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Post-herpetic neuralgia: Diagnosis of post-herpetic neuralgia
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

LIVTENCITY

Products Affected

- Livtencity

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of cytomegalovirus (CMV) infection/disease. Patient is a recipient of one of the following: a) Hematopoietic stem cell transplant OR b) Solid organ transplant. Trial and failure of a minimum 2 weeks duration, contraindication, or intolerance to one of the following therapies at an appropriately indicated dose: a) Intravenous (IV) ganciclovir, b) Oral valganciclovir, c) IV foscarnet, OR d) IV cidofovir. Patient weighs greater than or equal to 35 kg.
Age Restrictions	Patient is 12 years of age or older.
Prescriber Restrictions	Prescribed by or in consultation with a provider who specializes in one of the following areas: transplant, oncology, or infectious disease.
Coverage Duration	8 weeks
Other Criteria	N/A

LONSURF

Products Affected

- Lonsurf

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Metastatic colorectal cancer (mCRC): Diagnosis of mCRC. Trial and failure, contraindication, or intolerance with all of the following: fluoropyrimidine-based chemotherapy, oxaliplatin-based chemotherapy, irinotecan-based chemotherapy, and anti-VEGF biological therapy (e.g., bevacizumab). One of the following: a) tumor is RAS mutant-type or b) tumor is RAS wild-type and Trial and failure, contraindication or intolerance to one anti-EGFR therapy (eg, Vectibix [panitumumab], Erbitux [cetuximab]). Gastric/Gastroesophageal Junction Adenocarcinoma: Diagnosis of metastatic gastric cancer or diagnosis of metastatic gastroesophageal junction adenocarcinoma. Trial and failure, contraindication or intolerance to at least two of the following: fluopyrimidine-based chemotherapy (e.g. fluorouracil), Platinum-based chemotherapy (e.g., carboplatin, cisplatin, oxaliplatin), Taxane (e.g., docetaxel, paclitaxel) or irinotecan-based chemotherapy, HER2/neu-targeted therapy (e.g., trastuzumab) (if HER2 overexpression).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LORBRENA

Products Affected

- Lorbrena

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. One of the following: A) Disease is advanced, metastatic, or recurrent and anaplastic lymphoma kinase (ALK)-positive OR B) Both of the following: 1) Disease is both of the following: i) advanced, metastatic, or recurrent and ii) ROS proto-oncogene 1 (ROS1)-positive AND 2) Disease has progressed on at least one of the following therapies: Xalkori (crizotinib), Rozlytrek (entrectinib), or Zykadia (ceritinib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LUMAKRAS

Products Affected

- Lumakras TABS 120MG, 320MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is one of the following: a) recurrent, b) advanced or c) metastatic. Tumor is KRAS G12C-mutated. Patient has received at least one prior systemic therapy (e.g., immune checkpoint inhibitor, platinum-based chemotherapy).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

LUMRYZ

Products Affected

- Lumryz

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Narcolepsy with cataplexy (Narcolepsy Type 1)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are present, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present. Narcolepsy without cataplexy (Narcolepsy Type 2)(initial): Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible), AND symptoms of cataplexy are absent, AND symptoms of excessive daytime sleepiness (eg, irrepressible need to sleep or daytime lapses into sleep) are present, AND Trial and failure, contraindication, or intolerance to both of the following: 1) modafinil, AND 2) methylphenidate-based stimulant.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 6 months. Reauth: plan year
Other Criteria	Narcolepsy Type 1 (reauth): Documentation demonstrating a reduction in the frequency of cataplexy attacks associated with therapy, OR documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy. Narcolepsy Type 2 (reauth): Documentation demonstrating a reduction in symptoms of excessive daytime sleepiness associated with therapy.

LUPRON DEPOT

Products Affected

- Lupron Depot (1-month)
- Lupron Depot (3-month)
- Lupron Depot (4-month)
- Lupron Depot (6-month)

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer (7.5 mg, 22.5 mg, 30 mg, 45 mg): Treatment of advanced prostate cancer. Endometriosis (3.75 mg, 11.25 mg) (initial): Diagnosis of endometriosis. One of the following: a) Patient has had surgical ablation to prevent recurrence, or b) trial and failure, contraindication, or intolerance to one NSAID (e.g., diclofenac, ibuprofen, meloxicam, naproxen) and one oral contraceptive (e.g., norethindrone-ethinyl estradiol, estradiol and norethindrone). Endometriosis (3.75 mg, 11.25 mg) (reauthorization): Symptoms recur after one course. Used in combination with one of the following: norethindrone 5 mg daily, other "add -back" sex hormones (e.g., estrogen, medroxyprogesterone), or other bone-sparing agents (e.g., bisphosphonates such as alendronate, risedronate). Uterine Leiomyomata (3.75 mg, 11.25 mg) (fibroids): Either for use prior to surgery to reduce size of fibroids to facilitate a surgical procedure (eg, myomectomy, hysterectomy) Or all of the following: treatment of anemia, anemia caused by uterine leiomyomata (fibroids), and use prior to surgery.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prostate CA: plan yr. Endometrosis(all), Uterine leiomyomata (anemia): 6 mo. (fibroids): 4 mo.
Other Criteria	Prostate Cancer (7.5 mg, 22.5 mg, 30 mg, 45 mg): Approve for continuation of prior therapy if within the past 120 days.

LUPRON DEPOT PED

Products Affected

- Lupron Depot-ped (1-month) INJ
7.5MG
- Lupron Depot-ped (3-month) INJ
11.25MG
- Lupron Depot-ped (6-month)

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Central Precocious Puberty (CPP) (initial): diagnosis of central precocious puberty (idiopathic or neurogenic). Onset of secondary sexual characteristics in one of the following: females less than age 8 or males less than age 9. Confirmation of diagnosis defined by one of the following: pubertal basal level of luteinizing hormone (based on laboratory reference ranges), a pubertal response to a GnRH stimulation test, or bone age advanced one year beyond the chronological age.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	CPP (ini, reauth): plan yr.
Other Criteria	CPP (Reauthorization): Documentation of bone age monitoring (eg, radiographic imaging).

LYNPARZA (TABLETS)

Products Affected

- Lynparza TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>High risk early breast cancer: Diagnosis of high risk early breast cancer. Presence of deleterious or suspected deleterious germline BRCA-mutations. Disease is human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) Disease is hormone receptor (HR)-negative, OR b) Both of the following: i) Disease is HR-positive AND ii) Patient is continuing concurrent treatment with endocrine therapy. Patient has been previously treated with neoadjuvant or adjuvant chemotherapy (e.g., anthracycline, taxane). Metastatic or recurrent breast cancer: Diagnosis of breast cancer. Disease is metastatic or recurrent. Presence of deleterious or suspected deleterious germline BRCA-mutations. Disease is human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) Disease is hormone receptor (HR) negative, or b) Disease is hormone receptor (HR)-positive and one of the following: i) disease has progressed on previous endocrine therapy or ii) provider attestation that treatment with endocrine therapy is inappropriate for the patient's disease. Pancreatic adenocarcinoma: Diagnosis of pancreatic adenocarcinoma. Disease is metastatic. Presence of deleterious or suspected deleterious germline BRCA-mutations. Disease has not progressed while receiving at least 16 weeks of a first-line platinum-based chemotherapy regimen (e.g., FOLFIRINOX, FOLFOX, etc.).</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Ovarian cancer (Maintenance Therapy): Diagnosis of one of the following: epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. Disease is one of the following: advanced or recurrent. One of the following: 1) Patient has had a complete or partial response to platinum-based chemotherapy (e.g., carboplatin, cisplatin), or 2) Both of the following: a) patient has had a complete or partial response to first-line platinum-based chemotherapy (e.g., carboplatin, cisplatin) AND b) one of the following: i) presence of deleterious or suspected deleterious

germline or somatic BRCA-mutations OR ii) both of the following: cancer is associated with homologous recombination deficiency (HRD)-positive status defined by either a deleterious or suspected deleterious BRCA mutation or genomic instability AND used in combination with bevacizumab (e.g., Avastin, Myvasi). Will be used as maintenance therapy. Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. One of the following: 1) Both of the following: a) Presence of deleterious or suspected deleterious homologous recombination repair (HRR) gene mutations and b) Disease has progressed following prior treatment with one of the following: i) enzalutamide (Xtandi), or ii) abiraterone (e.g., Zytiga, Yonsa), OR 2) All of the following: a) Presence of deleterious or suspected deleterious BRCA-mutation, b) Used in combination with abiraterone (e.g., Zytiga, Yonsa), and c) Used in combination with prednisone or prednisolone. One of the following: 1) used in combination with a gonadotropin-releasing hormone (GnRH) analog [e.g., Lupron (leuprolide), Zoladex (goserelin), Trelstar (triptorelin), Vantas (histrelin), Firmagon (degarelix)], or 2) Patient has had bilateral orchiectomy. All indications: Approve for continuation of prior therapy if within the past 120 days.

LYTGOBI

Products Affected

- Lytgobi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of intrahepatic cholangiocarcinoma. Disease is one of the following: a) unresectable locally advanced, or b) metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangements. Patient has been previously treated.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

MAVYRET

Products Affected

- Mavyret

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine
Coverage Duration	8 to 16 weeks. Criteria applied consistent with current AASLD/IDSA guideline
Other Criteria	N/A

MEKINIST

Products Affected

- Mekinist

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable or metastatic melanoma AND cancer is BRAF V600 mutant type. Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type. Involvement of lymph nodes following complete resection. Used as adjuvant therapy. Medication is used in combination with Tafinlar (dabrafenib). Non-small Cell Lung Cancer (NSCLC): All of the following: diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type AND medication is used in combination with Tafinlar (dabrafenib). Anaplastic Thyroid Cancer (ATC): Diagnosis of ATC. One of the following: 1) Disease is one of the following: metastatic, locally advanced, or unresectable OR 2) Prescribed as adjuvant therapy following resection. Cancer is BRAF V600E mutant type. Medication is used in combination with Tafinlar (dabrafenib). Solid tumors: Presence of solid tumor. Disease is unresectable or metastatic. Patient has progressed on or following prior systemic treatment (e.g., carboplatin, 5-fluorouracil, paclitaxel). Cancer is BRAF V600E mutant type. Medication is used in combination with Tafinlar (dabrafenib). Low-grade glioma: Diagnosis of low-grade glioma. Cancer is BRAF V600E mutant type. Medication is used in combination with Tafinlar (dabrafenib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

MEKTOVI

Products Affected

- Mektovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma. Patient is positive for BRAF V600 mutation. Used in combination with Braftovi (encorafenib). Non-Small Cell Lung Cancer (NSCLC): Diagnosis of metastatic NSCLC. Patient is positive for BRAF V600 mutation. Used in combination with Braftovi (encorafenib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

MEMANTINE

Products Affected

- Memantine Hcl Titration Pak
- Memantine Hydrochloride SOLN
2MG/ML
- Memantine Hydrochloride TABS
- Memantine Hydrochloride Er
- Namzaric

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Age 41 or older, or diagnosis of moderate to severe dementia of the Alzheimer's type.
Age Restrictions	No Prior Authorization if patient is age 41 or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

MENACTRA

Products Affected

- Menactra

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroups A, C, W, and Y. Patient has not received Menactra, Menveo, or Menquadfi vaccine in the previous 12 months. One of the following: 1) Age lesser than or equal to 55 years, or 2) Patient is in a high risk group as defined by the Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months (2 vaccines per year)
Other Criteria	N/A

MENQUADFI

Products Affected

- Menquadfi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroups A, C, W, and Y. Patient has not received Menactra, Menveo, or Menquadfi vaccine in the previous 12 months. One of the following: 1) Age greater than or equal to 2 years, or 2) Patient is in a high risk group as defined by the Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month (1 vaccine per year)
Other Criteria	N/A

MENVEO

Products Affected

- Menveo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroups A, C, W, and Y. Patient has not received Menactra, Menveo, or Menquadfi vaccine in the previous 12 months. One of the following: 1) Age lesser than or equal to 55 years, or 2) Patient is in a high risk group as defined by the Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year (4 vaccines per year)
Other Criteria	N/A

METHOTREXATE INJECTION PREFERRED

Products Affected

- Rasuvo INJ 10MG/0.2ML, 12.5MG/0.25ML, 15MG/0.3ML, 17.5MG/0.35ML, 20MG/0.4ML, 22.5MG/0.45ML, 25MG/0.5ML, 30MG/0.6ML, 7.5MG/0.15ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid arthritis (RA): Diagnosis of severe, active RA. Polyarticular juvenile idiopathic arthritis (PJIA): Diagnosis of active PJIA. Psoriasis: Diagnosis of severe psoriasis.
Age Restrictions	N/A
Prescriber Restrictions	RA, PJIA: Prescribed by or in consultation with a rheumatologist. Psoriasis: Prescribed by or in consultation with a dermatologist.
Coverage Duration	Plan year
Other Criteria	N/A

MIGLUSTAT

Products Affected

- Miglustat
- Yargesa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gaucher disease: Diagnosis of mild to moderate Type 1 Gaucher disease. Patient is unable to receive enzyme replacement therapy due to one of the following conditions: allergy or hypersensitivity to enzyme replacement therapy, poor venous access, or unavailability of enzyme replacement therapy (e.g. Cerezyme, VPRIV).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

MODAFINIL

Products Affected

- Modafinil TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Obstructive sleep apnea (OSA) (Initial): Diagnosis (dx) of OSA defined by one of the following: 15 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), or both of the following: 5 or more obstructive respiratory events per hour of sleep confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible), and 1 of the following symptoms: unintentional sleep episodes during wakefulness, daytime sleepiness, unrefreshing sleep, fatigue, insomnia, waking up breath holding/gasping/choking, loud snoring, or breathing interruptions during sleep. Shift-work disorder (SWD) (Initial):Dx of SWD confirmed by one of the following: 1) symptoms of excessive sleepiness or insomnia for at least 3 months, which is associated with a work period (usually night work) that occurs during the normal sleep period, OR 2) A sleep study demonstrating loss of a normal sleep-wake pattern (ie, disturbed chronobiologic rhythmicity). Confirmation that no other medical conditions or medications are causing the symptoms of excessive sleepiness or insomnia. Narcolepsy (initial): Dx of narcolepsy as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). MS Fatigue (initial): Dx of multiple sclerosis (MS). Patient is experiencing fatigue. Idiopathic Hypersomnia (initial): Dx of idiopathic hypersomnia as confirmed by a sleep study (unless prescriber provides justification confirming that a sleep study is not feasible). Depression (initial): Treatment-resistant depression defined as diagnosis of major depressive disorder (MDD) or bipolar depression, AND trial and failure, contraindication, or intolerance to at least two antidepressants from different classes (eg, SSRIs, SNRIs, bupropion). Used as adjunctive therapy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	OSA,SWD,MS Fatigue,Hypersomnia,Depression:Initial,Reauth:6 mo.Narcolepsy:Initial,Reauth:Plan Yr

Other Criteria	OSA, Narcolepsy, Idiopathic Hypersomnia (Reauth): Patient demonstrates positive clinical response to therapy. SWD (Reauth): Patient demonstrates positive clinical response to therapy. MS Fatigue (reauth): Patient is experiencing relief of fatigue with therapy. Depression (reauth): Patient demonstrates positive clinical response to therapy. Used as adjunctive therapy.
-----------------------	---

MOUNJARO

Products Affected

- Mounjaro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

MRESVIA

Products Affected

- Mresvia

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV). Patient has not received an RSV vaccine (i.e., Abrysvo, Arexvy) in their lifetime. One of the following: a) Age greater than or equal to 75 years, or b) Both of the following: 1) Age 60 through 74 years, and 2) Patient is at increased risk of severe RSV disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

NAYZILAM

Products Affected

- Nayzilam

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Epilepsy: Diagnosis of epilepsy. Frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

NERLYNX

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Early Stage Breast cancer: Diagnosis (dx) of early stage breast cancer. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Patient has received adjuvant trastuzumab based therapy (e.g., Herceptin, Kanjinti, etc.). Advanced or Metastatic Breast Cancer: 1) All of the following: a) Dx of advanced or metastatic breast cancer, b) Disease is human epidermal growth factor receptor 2 (HER2)-positive, c) Patient has received two or more prior anti-HER2 based regimens (e.g., trastuzumab + pertuzumab + docetaxel, ado-trastuzumab emtansine, etc.), and d) Used in combination with capecitabine, OR 2) Both of the following: a) Diagnosis of Stage IV (M1) breast cancer, and b) One of the following: i) Hormone receptor-positive, (HER2)-negative disease in patients who have already received a CDK4/6 inhibitor therapy, or ii) Triple negative disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

NEXLETOL

Products Affected

- Nexletol

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Heterozygous familial hypercholesterolemia (HeFH) or primary hyperlipidemia (Initial): One of the following diagnoses: A) Heterozygous familial hypercholesterolemia (HeFH), OR B) Primary hyperlipidemia. One of the following: (1) used as adjunct to statin therapy, (2) patient has intolerance to a statin, OR (3) patient has a contraindication to all statins. One of the following LDL-C values within the last 120 days: (1) LDL-C greater than or equal to 55 mg/dL with ASCVD or (2) LDL-C greater than or equal to 70 mg/dL without ASCVD. Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD (Initial): One of the following diagnoses: A) Established CVD (e.g., coronary artery disease, symptomatic peripheral arterial disease, cerebrovascular atherosclerotic disease), OR B) a high risk for a CVD event but without established CVD [e.g., diabetes mellitus (type 1 or type 2) in females over 65 years of age or males over 60 years of age]. One of the following: (1) used as adjunct to statin therapy, (2) patient has intolerance to a statin, OR (3) patient has a contraindication to all statins. One of the following LDL-C values within the last 120 days: (1) LDL-C greater than or equal to 55 mg/dL with ASCVD or (2) LDL-C greater than or equal to 70 mg/dL without ASCVD.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 6 months. Reauth: Plan year
Other Criteria	HeFH or primary hyperlipidemia (Reauth): Patient demonstrates positive clinical response to therapy. Patient continues to receive other lipid-lowering treatment (e.g., statins, ezetimibe) or patient has an inability to take other lipid-lowering therapy (e.g., statins, ezetimibe). Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD (Reauth): Patient demonstrates positive clinical response to therapy.

NEXLIZET

Products Affected

- Nexlizet

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Heterozygous familial hypercholesterolemia (HeFH) or primary hyperlipidemia (Initial): One of the following diagnoses: A) Heterozygous familial hypercholesterolemia (HeFH), OR B) Primary hyperlipidemia. One of the following: (1) used as adjunct to statin therapy, (2) patient has intolerance to a statin, OR (3) patient has a contraindication to all statins. One of the following LDL-C values within the last 120 days: (1) LDL-C greater than or equal to 55 mg/dL with ASCVD or (2) LDL-C greater than or equal to 70 mg/dL without ASCVD. Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD (Initial): One of the following diagnoses: A) Established CVD (e.g., coronary artery disease, symptomatic peripheral arterial disease, cerebrovascular atherosclerotic disease), OR B) a high risk for a CVD event but without established CVD [e.g., diabetes mellitus (type 1 or type 2) in females over 65 years of age or males over 60 years of age]. One of the following: (1) used as adjunct to statin therapy, (2) patient has intolerance to a statin, OR (3) patient has a contraindication to all statins. One of the following LDL-C values within the last 120 days: (1) LDL-C greater than or equal to 55 mg/dL with ASCVD or (2) LDL-C greater than or equal to 70 mg/dL without ASCVD.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 6 months. Reauth: Plan year
Other Criteria	HeFH or primary hyperlipidemia (Reauth): Patient demonstrates positive clinical response to therapy. Patient continues to receive other lipid-lowering treatment (e.g., statins, ezetimibe) or patient has an inability to take other lipid-lowering therapy (e.g., statins, ezetimibe). Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD (Reauth): Patient demonstrates positive clinical response to therapy.

NINLARO

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple myeloma: Diagnosis of multiple myeloma. One of the following: 1) Both of the following: Patient has received at least one prior therapy for multiple myeloma [e.g., Velcade (bortezomib)] AND Used as part of combination regimen including dexamethasone [combination regimen may include additional agents, such as Revlimid (lenalidomide)] OR 2) Both of the following: a) Used as primary therapy or maintenance therapy and b) Patient is not a transplant candidate OR 3) Both of the following: a) Patient is a transplant candidate and b) One of the following: Patient has symptomatic disease following response to primary myeloma therapy or response or stable disease following autologous or allogeneic stem cell transplant.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

NUBEQA

Products Affected

- Nubeqa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Non-metastatic castration-resistant or castration-recurrent prostate cancer (NM-CRPC): Diagnosis of non-metastatic castration-resistant (chemical or surgical) or castration-recurrent prostate cancer. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., Zoladex [goserelin], Vantas [histrelin], Lupron [leuprolide], Trelstar [triptorelin]) OR 2) Patient received bilateral orchiectomy.</p> <p>Metastatic hormone-sensitive prostate cancer (mHSPC): Diagnosis of mHSPC. Used in combination with docetaxel. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g., Zoladex [goserelin], Vantas [histrelin], Lupron [leuprolide], Trelstar [triptorelin]) OR 2) Patient received bilateral orchiectomy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	NM-CRPC, mHSPC: Plan year
Other Criteria	NM-CRPC, mHSPC: Approve for continuation of prior therapy if within the past 120 days.

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pseudobulbar affect (PBA) (initial): Diagnosis of PBA, confirmed by one of the following: 1) Physician attestation that a baseline Center for Neurologic Studies Lability Scale (CNS-LS) score has been assessed OR 2) Patient attestation that patient has experienced involuntary, sudden, or frequent episodes of laughing and/or crying. Patient has a brain injury or neurologic disease from one of the following: amyotrophic lateral sclerosis, multiple sclerosis, Parkinson's disease, stroke, or traumatic brain injury. Patient does not have any of the following contraindications: a) Concomitant use with other drugs containing quinidine, quinine, or mefloquine, b) History of Nuedexta, quinine, mefloquine or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome, c) Known hypersensitivity to dextromethorphan (e.g., rash, hives), d) Taking monoamine oxidase inhibitors (MAOIs) (e.g., phenelzine, selegiline, tranylcypromine) or have taken MAOIs within the preceding 14 days, e) Has prolonged QT interval, congenital long QT syndrome or a history suggestive of torsades de pointes, or has heart failure, f) Receiving drugs that both prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), g) Has complete atrioventricular (AV) block without implanted pacemakers, or at high risk of complete AV block.
Age Restrictions	N/A
Prescriber Restrictions	PBA (initial/reauth): Prescribed by or in consultation with one of the following specialists: neurologist, psychiatrist.
Coverage Duration	PBA (initial/reauth): plan year
Other Criteria	PBA (reauth): One of the following: 1) Physician attestation that the patient's CNS-LS score has improved since baseline OR 2) Physician attestation that frequency of laughing and/or crying episodes has decreased since baseline. Diagnosis of PBA. Patient has a brain injury or neurologic disease from one of the following: amyotrophic lateral sclerosis, multiple sclerosis, Parkinson's disease, stroke, or traumatic brain injury. Patient does not have any of the following contraindications: a) Concomitant use with other drugs containing quinidine, quinine, or

	mefloquine, b) History of Nuedexta, quinine, mefloquine or quinidine-induced thrombocytopenia, hepatitis, bone marrow depression, or lupus-like syndrome, c) Known hypersensitivity to dextromethorphan (e.g., rash, hives), d) Taking monoamine oxidase inhibitors (MAOIs) (e.g., phenelzine, selegiline, tranylcypromine) or have taken MAOIs within the preceding 14 days, e) Has prolonged QT interval, congenital long QT syndrome or a history suggestive of torsades de pointes, or has heart failure, f) Receiving drugs that both prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), g) Has complete atrioventricular (AV) block without implanted pacemakers, or at high risk of complete AV block.
--	--

NUPLAZID

Products Affected

- Nuplazid CAPS
- Nuplazid TABS 10MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Parkinson's disease psychosis: Diagnosis of Parkinson's disease. Patient has at least one of the following: hallucinations or delusions.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

NURTEC

Products Affected

- Nurtec

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Treatment of Migraine (initial): Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. Medication will not be used in combination with another CGRP inhibitor for the acute treatment of migraines. Preventive Treatment of Episodic Migraine (EM) (initial): Diagnosis of EM. Patient has greater than or equal to 4 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) Topiramate (Topamax), e) Venlafaxine (Effexor), f) Candesartan (Atacand). Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All Indications (initial, reauth): Plan year
Other Criteria	Acute Treatment of Migraine (reauth): Patient demonstrates positive clinical response to therapy. Medication will not be used in combination with another CGRP inhibitor for the acute treatment of migraines. Preventive Treatment of EM (reauth): Patient demonstrates positive clinical response to therapy. Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.

OCTREOTIDE

Products Affected

- Octreotide Acetate INJ
1000MCG/ML, 100MCG/ML,
200MCG/ML, 500MCG/ML,
50MCG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Acromegaly (initial): Diagnosis of acromegaly confirmed by one of the following: serum GH level greater than 1 ng/mL after a 2-hour oral glucose tolerance test at the time of diagnosis, or elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at the time of diagnosis. One of the following: A) Inadequate response to surgery, radiotherapy, or dopamine agonist (e.g., bromocriptine, cabergoline) therapy, or B) Not a candidate for any of the following: surgery, radiotherapy, dopamine agonist (e.g., bromocriptine, cabergoline) therapy. HIV/AIDS-Related Diarrhea (initial): Diagnosis of HIV/AIDS-related diarrhea. Carcinoid tumors, symptomatic treatment of diarrhea or flushing (initial): diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes. Vasoactive Intestinal Peptide Tumors, symptomatic treatment of diarrhea (initial): Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea. Cancer Chemotherapy- and/or Radiation- Induced Diarrhea (initial): Diagnosis of complicated diarrhea due to concurrent cancer chemotherapy and/or radiation or uncomplicated diarrhea due to concurrent cancer chemotherapy and/or radiation. Carcinoid tumor: diagnosis of carcinoid tumor. Reauthorization (all except carcinoid tumor): Patient demonstrates positive clinical response to therapy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All Indications: Plan Year
Other Criteria	<p>Uncomplicated diarrhea due to concurrent cancer chemotherapy and/or radiation (initial): Trial and failure, contraindication, or intolerance (TF/C/I) to standard therapy (e.g., loperamide). HIV/AIDS-related Diarrhea (initial): TF/C/I to standard therapy (e.g., loperamide, diphenoxylate with atropine). Carcinoid tumor: Approve for continuation of prior therapy if within the past 120 days.</p>

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced basal cell carcinoma: Diagnosis of locally advanced basal cell carcinoma. One of the following: cancer that has recurred following surgery or radiation therapy, or patient is not a candidate for surgery or radiation therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

OFEV

Products Affected

- Ofev

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF, defined as exclusion of other known causes of interstitial lung disease and either the presence of usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF in patients not subjected to lung biopsy, or HRCT and surgical lung biopsy pattern revealing IPF or probable IPF in patients subjected to a lung biopsy.</p> <p>Systemic Sclerosis-associated interstitial lung disease (SSc-ILD) (initial): Diagnosis of SSc-ILD, defined as exclusion of other known causes of interstitial lung disease (ILD) and either the presence of idiopathic interstitial pneumonia (e.g., fibrotic nonspecific interstitial pneumonia [NSIP], usual interstitial pneumonia [UIP] and centrilobular fibrosis) pattern on HRCT revealing SSc-ILD or probable SSc-ILD in patients not subjected to surgical lung biopsy, or HRCT and surgical lung biopsy pattern revealing SSc-ILD or probable SSc-ILD in patients subjected to a lung biopsy.</p> <p>Chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype (initial): Diagnosis of chronic fibrosing ILDs. Patient has a high-resolution computed tomography (HRCT) showing at least 10% of lung volume with fibrotic features. Disease has a progressive phenotype as observed by one of the following: decline of forced vital capacity (FVC), worsening of respiratory symptoms, or increased extent of fibrosis seen on imaging.</p> <p>IPF, SSc-ILD, Chronic Fibrosing ILDs with a progressive phenotype (reauth): Patient demonstrates positive clinical response to therapy.</p>
Age Restrictions	N/A
Prescriber Restrictions	(initial): Prescribed by or in consultation with a pulmonologist.
Coverage Duration	(initial, reauth): plan year
Other Criteria	N/A

OGSIVEO

Products Affected

- Ogsiveo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of desmoid tumor. Patient requires systemic treatment. Disease is progressive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

OJEMDA

Products Affected

- Ojemda

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of pediatric low-grade glioma. Disease is relapsed or refractory. Disease has a BRAF fusion or rearrangement, or BRAF V600 mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

OJJAARA

Products Affected

- Ojjaara

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of ONE of the following: a) Primary myelofibrosis, b) Post-polycythemia vera myelofibrosis, OR c) Post-essential thrombocythemia myelofibrosis. Disease is intermediate or high risk. Patient has anemia.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ONUREG

Products Affected

- Onureg

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of acute myeloid leukemia (AML). Patient has received previous treatment with an intensive induction chemotherapy regimen (e.g., cytarabine + daunorubicin, cytarabine + idarubicin). Patient has achieved one of the following: a) first complete remission (CR) or b) complete remission with incomplete blood count recovery (CRi). Patient is not a candidate for intensive curative therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

OPSUMIT

Products Affected

- Opsumit

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH: Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: plan year
Other Criteria	N/A

ORENCIA SC

Products Affected

- Orenzia INJ 125MG/ML,
50MG/0.4ML, 87.5MG/0.7ML
- Orenzia Clickject

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate (MTX), leflunomide, sulfasalazine. TF/C/I to one of the following: Enbrel (etanercept), one Formulary adalimumab product, Rinvoq (upadacitinib), or Xeljanz/Xeljanz XR (tofacitinib), OR for continuation of prior therapy if within the past 120 days. Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of moderately to severely active PJIA. Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. One of the following: TF/C/I to Enbrel (etanercept), one Formulary adalimumab product, Xeljanz (tofacitinib), OR for continuation of prior therapy if within the past 120 days. Psoriatic Arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. TF/C/I to one of the following: Enbrel (etanercept), one Formulary adalimumab product, Cosentyx SC (secukinumab), Stelara (ustekinumab), Skyrizi (risankizumab-rzaa), Xeljanz/Xeljanz XR (tofacitinib), Rinvoq (upadacitinib), or Otezla (apremilast), OR for continuation of prior therapy if within the past 120 days.
Age Restrictions	N/A
Prescriber Restrictions	RA (initial), JIA (initial): Prescribed by or in consultation with a rheumatologist. PsA (Initial): Prescribed by or in consultation with a dermatologist or rheumatologist.
Coverage Duration	All indications (Initial): 6 months, (Reauth): plan year
Other Criteria	All indications (Reauth): Patient demonstrates positive clinical response to therapy.

ORGOVYX

Products Affected

- Orgovyx

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ORKAMBI

Products Affected

- Orkambi TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic fibrosis (initial): Diagnosis of cystic fibrosis (CF). Submission of laboratory records confirming the patient is homozygous for the F508del mutation in the CFTR gene.
Age Restrictions	Patient is greater than or equal to 6 years of age
Prescriber Restrictions	CF (initial, reauthorization): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
Coverage Duration	Initial: 6 months. Reauth: plan year
Other Criteria	CF (reauth): Patient demonstrates positive clinical response to therapy.

ORKAMBI GRANULES

Products Affected

- Orkambi PACK

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cystic fibrosis (initial): Diagnosis of cystic fibrosis (CF). Submission of laboratory records confirming the patient is homozygous for the F508del mutation in the CFTR gene. One of the following: A) Patient is 1 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.
Age Restrictions	N/A
Prescriber Restrictions	CF (initial, reauthorization): Prescribed by or in consultation with a specialist affiliated with a CF care center or pulmonologist
Coverage Duration	Initial: 6 months. Reauth: plan year
Other Criteria	CF (Reauthorization): Patient demonstrates positive clinical response to therapy. One of the following: A) Patient is 1 through 5 years of age, OR B) Both of the following: Patient is 6 years of age or greater AND Patient is unable to swallow oral tablets.

ORSERDU

Products Affected

- Orserdu

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of breast cancer. Disease is advanced or metastatic. One of the following: a) Patient is male, b) Patient is a postmenopausal woman, or c) Patient is a premenopausal woman treated with ovarian ablation/suppression. Disease is estrogen receptor (ER)-positive. Disease is human epidermal growth factor receptor 2 (HER2)-negative. Presence of estrogen receptor (ESR1) mutation(s). Disease has progressed following at least one line of endocrine therapy [e.g., Faslodex (fulvestrant)].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

OSPHENA

Products Affected

- Ospheña

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Dyspareunia: Diagnosis of moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause. Vaginal dryness: Diagnosis of moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

OTEZLA

Products Affected

- Otezla

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Psoriatic arthritis (PsA, initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Plaque psoriasis (initial): Diagnosis of plaque psoriasis. Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Oral ulcers associated with Behcet’s Disease (initial): Diagnosis of Behcet’s Disease. Patient has active oral ulcers.
Age Restrictions	N/A
Prescriber Restrictions	PsA (init): Prescribed by or in consultation with one of the following specialists: dermatologist or rheumatologist. Plaque psoriasis (init): Prescribed by or in consultation with a dermatologist.
Coverage Duration	All uses (initial): 6 months. All uses (reauth): plan year.
Other Criteria	All indications (Reauth): Patient demonstrates positive clinical response to therapy.

OZEMPIC

Products Affected

- Ozempic INJ 2MG/3ML, 4MG/3ML, 8MG/3ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM. Nonalcoholic steatohepatitis (NASH)/Metabolic dysfunction-associated steatohepatitis (MASH) (Initial): Diagnosis of MASH, formerly known as nonalcoholic steatohepatitis (NASH). Patient does not have cirrhosis (e.g., decompensated cirrhosis). Submission of medical records (e.g., chart notes) confirming diagnosis has been confirmed by one of the following: FibroScan-aspartate aminotransferase (FAST), MRI-aspartate aminotransferase (MAST), or liver biopsy. Submission of medical records (e.g., chart notes) confirming disease is fibrosis stage F2 or F3 as confirmed by one of the following: FibroScan, Fibrosis-4 index (FIB-4), or Magnetic Resonance Elastography (MRE).
Age Restrictions	N/A
Prescriber Restrictions	MASH (Initial): Prescribed by or in consultation with a gastroenterologist or hepatologist.
Coverage Duration	Plan year
Other Criteria	MASH (Reauth): Patient demonstrates positive response to therapy.

PANRETIN

Products Affected

- Panretin

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Kaposi's sarcoma lesions: Diagnosis of cutaneous lesions in patients with AIDS-related Kaposi's sarcoma (KS). Not used when systemic anti-KS therapy is required (e.g., more than 10 new KS lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary KS, or symptomatic visceral involvement).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

PEGFILGRASTIM PREFERRED

Products Affected

- Neulasta
- Udenyca

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Neutropenia Associated with Dose Dense Chemotherapy (NDDC): Patient is receiving National Cancer Institute's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer, or a dose-dense chemotherapy regimen for which the incidence of febrile neutropenia is unknown. Chemotherapy-Induced Febrile Neutropenia (CFN): Patient is receiving a chemotherapy regimen associated with greater than 20% incidence of febrile neutropenia, or patient is receiving chemotherapy regimen associated with 10-20% incidence of febrile neutropenia and has 1 or more risk factors associated with chemotherapy-induced infection, febrile neutropenia or neutropenia. Secondary prophylaxis of FN: For patients who are receiving myelosuppressive anticancer drugs associated with neutropenia. Patient has a history of FN or dose-limiting event during a previous course of chemotherapy (secondary prophylaxis). Acute radiation syndrome (ARS): Patient was/will be acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of ARS).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	ARS: 1 mo. CFN, NDDC, FN (prophylaxis): 3 mo or duration of tx.
Other Criteria	N/A

PEGINTERFERON ALFA - 2A

Products Affected

- Pegasys

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Hepatitis B: Diagnosis of chronic hepatitis B infection. Chronic Hepatitis C: Diagnosis of chronic hepatitis C infection. Patient has compensated liver disease. One of the following: a) Used in combination with one other hepatitis C virus (HCV) antiviral drug (e.g., Mavyret [glecaprevir-pibrentasvir], ribavirin) OR b) Both of the following: Used as monotherapy AND contraindication or intolerance to all other HCV antiviral drugs (e.g., Mavyret [glecaprevir-pibrentasvir], ribavirin).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HepB, HepC: 48 wks.
Other Criteria	N/A

PEMAZYRE

Products Affected

- Pemazyre

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cholangiocarcinoma: Diagnosis of cholangiocarcinoma. Disease is one of the following: unresectable locally advanced or metastatic. Disease has presence of a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement. Patient has been previously treated. Myeloid/Lymphoid neoplasms: Diagnosis of myeloid/lymphoid neoplasm (MLNs). Disease is relapsed or refractory. Disease has presence of a fibroblast growth factor receptor 1 (FGFR1) rearrangement.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

PENBRAYA

Products Affected

- Penbraya

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroups A, B, C, W, and Y. Patient has not received or will not receive more than two Penbraya vaccines in the previous 12 months. Age lesser than or equal to 25 years.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year (2 vaccinations per year)
Other Criteria	N/A

PIQRAY

Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast Cancer (BC): Diagnosis of advanced or metastatic BC. Disease is hormone receptor (HR)-positive, and human epidermal growth factor receptor 2 (HER2)-negative. Presence of one or more PIK3CA mutations. Used in combination with fulvestrant. Disease has progressed on or after an endocrine-based regimen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

PIRFENIDONE

Products Affected

- Pirfenidone

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Idiopathic pulmonary fibrosis (IPF) (initial): Diagnosis of IPF, defined as exclusion of other known causes of interstitial lung disease and either the presence of usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF in patients not subjected to lung biopsy, or HRCT and surgical lung biopsy pattern revealing IPF or probable IPF in patients subjected to a lung biopsy. IPF (reauth): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	(initial): Prescribed by or in consultation with a pulmonologist.
Coverage Duration	(initial, reauth): plan year
Other Criteria	N/A

POMALYST

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple myeloma: Diagnosis of multiple myeloma. Used in combination with dexamethasone. Trial and failure, contraindication or intolerance to both an immunomodulatory agent [eg, Revlimid (lenalidomide)] and a proteasome inhibitor [eg, Velcade (bortezomib)]. Kaposi sarcoma (KS): One of the following: 1) Both of the following: a) Diagnosis of AIDS-related KS and b) Patient has failed highly active antiretroviral therapy (HAART) [e.g., Biktarvy (bictegravir/emtricitabine/tenofovir alafenamide), Dovato (dolutegravir/lamivudine), Triumeq (dolutegravir/abacavir/lamivudine)], OR 2) Both of the following: a) Diagnosis of KS and b) Patient is HIV-negative.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

POSACONAZOLE

Products Affected

- Posaconazole Dr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prophylaxis of Invasive Fungal Infections (IFI): Used as prophylaxis of invasive fungal infections caused by <i>Aspergillus</i> or <i>Candida</i> for one of the following conditions: 1) Patient is at high risk of infections due to severe immunosuppression from hematopoietic stem cell transplant (HSCT) with graft-versus-host disease (GVHD) or hematologic malignancies with prolonged neutropenia from chemotherapy [eg, acute myeloid leukemia (AML), myelodysplastic syndrome (MDS)], OR 2) patient has a prior fungal infection requiring secondary prophylaxis. Treatment of IFI: Used as treatment of invasive fungal infections caused by <i>Aspergillus</i> .
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Prophylaxis of IFI: plan year. Treatment of IFI: 3 months.
Other Criteria	N/A

PREVYMIS ORAL

Products Affected

- Prevymis TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cytomegalovirus (CMV) Prophylaxis in Hematopoietic Stem Cell Transplant (HSCT): Used for prophylaxis of CMV infection and disease AND patient is a CMV-seropositive recipient [R+] of an allogeneic HSCT. CMV Prophylaxis in Kidney Transplant: Used for prophylaxis CMV infection and disease. Patient is a CMV-seronegative recipient [R-]. Patient is receiving a kidney transplant from a CMV-seropositive donor [D+].
Age Restrictions	N/A
Prescriber Restrictions	CMV Prophylaxis in HSCT: Prescribed by or in consultation with an oncologist, hematologist, physician experienced in the management of transplant patients, or infectious disease specialist. CMV Prophylaxis in Kidney Transplant: Prescribed by or in consultation with a nephrologist, physician experienced in the management of transplant patients, or infectious disease specialist.
Coverage Duration	CMV Prophylaxis in HSCT: 7 months. CMV Prophylaxis in Kidney Transplant: 7 months.
Other Criteria	N/A

PROMACTA

Products Affected

- Promacta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Idiopathic thrombocytopenic purpura (ITP) (initial): Diagnosis of one of the following: persistent ITP, chronic ITP or relapsed/refractory ITP. Baseline platelet count is less than 30,000/mcL. Patient's degree of thrombocytopenia and clinical condition increase the risk of bleeding. ITP (reauthorization): Patient demonstrates positive clinical response to therapy as evidenced by an increase in platelet count to a level sufficient to avoid clinically important bleeding. Chronic Hepatitis C-associated thrombocytopenia (initial): Diagnosis of chronic hepatitis C-associated thrombocytopenia. One of the following: planning to initiate and maintain interferon-based treatment or currently receiving interferon-based treatment. First-line for severe aplastic anemia (SAA): Diagnosis of SAA. Used for first-line treatment (i.e., patient has not received prior immunosuppressive therapy with any equine antithymocyte globulin plus cyclosporine, alemtuzumab, or high dose cyclophosphamide). Used in combination with standard immunosuppressive therapy (e.g., Atgam [antithymocyte globulin equine] and cyclosporine). Patient meets at least two of the following: 1) absolute neutrophil count less than 500/mcL, 2) platelet count less than 20,000/mcL, 3) absolute reticulocyte count less than 60,000/mcL. Refractory SAA (initial): Diagnosis of refractory SAA. Patient has a platelet count less than 30,000/mcL. SAA (reauthorization): Patient demonstrates positive clinical response to therapy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1stline SAA:6mo.HepC (init):3mo.RefractSAA(init):16wk.ITP,HepC(reauth),RefractSAA(reauth):plan yr
Other Criteria	ITP (initial): Trial and failure, contraindication, or intolerance to at least one of the following: corticosteroids (e.g., prednisone, methylprednisolone), immunoglobulins [e.g., Gammagard, immune globulin (human)], or splenectomy. Chronic Hepatitis C-associated thrombocytopenia

(Reauthorization): One of the following criteria: For patients that started treatment with eltrombopag prior to initiation of treatment with interferon, eltrombopag will be approved when both of the following are met: currently on antiviral interferon treatment for treatment of chronic hepatitis C and documentation that patient reached threshold platelet count that allows initiation of antiviral interferon therapy with eltrombopag treatment by week 9. OR for patients that started treatment with eltrombopag while on concomitant treatment with interferon, eltrombopag will be approved based on the following criterion: currently on antiviral interferon therapy for treatment of chronic hepatitis C. Refractory SAA: Trial and failure, contraindication, or intolerance to at least one course of immunosuppressive therapy (eg, Atgam (antithymocyte globulin equine), Thymoglobulin (antithymocyte globulin rabbit), cyclosporine).

PURIXAN

Products Affected

- Purixan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	One of the following: History of contraindication or intolerance to generic mercaptopurine tablets OR patient is unable to swallow tablets.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

PYRUKYND

Products Affected

- Pyrukynd
- Pyrukynd Taper Pack

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of hemolytic anemia confirmed by the presence of chronic hemolysis (e.g., increased indirect bilirubin, elevated lactated dehydrogenase [LDH], decreased haptoglobin, increased reticulocyte count). Diagnosis of pyruvate kinase deficiency confirmed by molecular testing of ALL the following mutations on the PKLR gene: a) Presence of at least 2 variant alleles in the pyruvate kinase liver and red blood cell (PKLR) gene, of which at least 1 was a missense variant AND b) Patients is not homozygous for the c.1436G to A (p.R479H) variant AND c) Patient does not have 2 non-missense variants (without the presence of another missense variant) in the PKLR gene. Hemoglobin is less than or equal to 10g/dL. Patient has symptomatic anemia or is transfusion dependent. Exclusion of other causes of hemolytic anemias (e.g., infections, toxins, drugs).
Age Restrictions	N/A
Prescriber Restrictions	Initial, Reauth: Prescribed by or in consultation with a hematologist.
Coverage Duration	Initial: 6 months. Reauth: Plan Year.
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy.

QINLOCK

Products Affected

- Qinlock

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gastrointestinal Stromal Tumor (GIST): Diagnosis of gastrointestinal stromal tumor (GIST). Disease is one of the following: a) advanced, b) metastatic, c) unresectable, or d) recurrent. One of the following: a) Trial and failure, contraindication, or intolerance to all of the following: imatinib (Gleevec), sunitinib (Sutent), and regorafenib (Stivarga), b) All of the following: performance status 0-2, history of progression on imatinib (Gleevec), and history of intolerance to sunitinib (Sutent), or c) All of the following: PDGFRA exon 18 mutations that are insensitive to imatinib (Gleevec) (including PDGFRA D842V), history of progression on avapritinib (Ayvakit), and history of progression on dasatinib (Sprycel).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

QUININE

Products Affected

- Quinine Sulfate CAPS 324MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis (dx) of uncomplicated malaria and one of the following: treatment in areas of chloroquine-sensitive malaria or treatment in areas of chloroquine-resistant malaria.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	7 days
Other Criteria	Chloroquine-sensitive malaria: Failure, contraindication or intolerance to chloroquine or hydroxychloroquine.

QULIPTA

Products Affected

- Qulipta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Episodic Migraine (EM) (initial): Diagnosis of EM. Patient has greater than or equal to 4 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) Topiramate (Topamax), e) Venlafaxine (Effexor), f) Candesartan (Atacand). Chronic Migraines (CM) (initial): Diagnosis of CM. Patient has greater than or equal to 8 migraine days per month. Trial and failure (after a trial of at least two months), contraindication, or intolerance to two of the following prophylactic therapies: a) Amitriptyline (Elavil), b) One of the following beta-blockers: atenolol, metoprolol, nadolol, propranolol, or timolol, c) Divalproex sodium (Depakote/Depakote ER), d) OnabotulinumtoxinA (Botox), e) Topiramate (Topamax), f) Venlafaxine (Effexor), g) Candesartan (Atacand). All Indications (initial): Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	EM, CM (initial, reauth): Plan year
Other Criteria	EM, CM (reauth): Patient demonstrates positive clinical response to therapy. Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines.

REGRANEX

Products Affected

- Regranex

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diabetic Neuropathic Ulcers: Patient has a lower extremity diabetic neuropathic ulcer. Treatment will be given in combination with ulcer wound care (eg, debridement, infection control, and/or pressure relief).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Diabetic Neuropathic Ulcers: 5 months.
Other Criteria	N/A

REPATHA

Products Affected

- Repatha
- Repatha Pushtronex System
- Repatha Sureclick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>HeFH/ASCVD/Primary HLD(init): One of the following diagnoses: A) Heterozygous familial hypercholesterolemia (HeFH), B) Atherosclerotic cardiovascular disease (ASCVD), OR C) Primary Hyperlipidemia (HLD). ONE of the following: A) One of the following LDL-C values while on maximally tolerated lipid lowering tx within the last 120 days: (1) LDL-C greater than or equal to 55 mg/dL with ASCVD. (2) LDL-C greater than or equal to 100 mg/dL without ASCVD. OR B) Both of the following: (1) Patient has been receiving PCSK9 therapy as adjunct to maximally tolerated lipid lowering therapy and (2) LDL-C values drawn within the past 12 months while on maximally tolerated lipid lowering therapy are within normal limits. AND One of the following: i) Pt has been receiving at least 12 weeks of one maximally-tolerated statin tx, ii) Pt is statin intolerant as evidenced by an inability to tolerate at least two statins, with at least one started at the lowest starting daily dose, due to intolerable symptoms or clinically significant biomarker changes of liver function or muscle function (e.g., creatine kinase), iii) patient has a contraindication to all statins. ONE of the following: Pt has been receiving ezetimibe (Zetia) tx as adjunct to max tolerated statin tx OR Pt has a hx of contraindication or intolerance to ezetimibe. HoFH (initial): dx of homozygous familial hypercholesterolemia (HoFH) as confirmed by one of the following: (1) Gen confirmation of 2 mutations in LDL receptor, ApoB, PCSK9, or LDLRAP1 or ARH, or (2) either untreated LDL greater than 500 or treated LDL greater than 300, AND either xanthoma before 10 yo or evidence of HeFH in both parents. Pt is receiving other lipid-lowering tx (eg statin, ezetimibe) OR patient has inability to take other lipid lowering tx (e.g., statin, ezetimibe).</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 6 months. Reauth: plan year
Other Criteria	HeFH/ASCVD/Primary HLD/HoFH (reauth): Pt continues to receive other lipid-lowering therapy (eg statin, ezetimibe) at the maximally

	tolerated dose or pt has an inability to take other lipid-lowering therapy (e.g., statin, ezetimibe). Patient demonstrates positive clinical response to therapy.
--	---

RETEVMO

Products Affected

- Retevmo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-Small Cell Lung cancer: Diagnosis of non-small cell lung cancer (NSCLC). Disease is one of the following: a) recurrent, b) advanced, or c) metastatic. Presence of RET gene fusion-positive or RET rearrangement positive tumor(s). Medullary Thyroid Cancer (MTC): Diagnosis of medullary thyroid cancer (MTC). Disease is advanced or metastatic. Disease has presence of RET gene mutation. Disease requires treatment with systemic therapy. Thyroid Cancer: Diagnosis of thyroid cancer. Disease is advanced or metastatic. Disease is RET gene fusion-positive. Disease requires treatment with systemic therapy. Patient is radioactive iodine-refractory or radioactive iodine therapy is not appropriate. Solid Tumors: Presence of RET gene fusion-positive solid tumor. Disease is recurrent, advanced, or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Non-Small Cell Lung Cancer, MTC, Thyroid Cancer, Solid Tumors: Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

REVCovi

Products Affected

- Revcovi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of adenosine deaminase deficiency (ADA) with severe combined immunodeficiency (SCID).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

REVLIMID

Products Affected

- Lenalidomide

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Myeloma: Diagnosis of multiple myeloma. Myelodysplastic syndrome (MDS) with a deletion 5q; Diagnosis of symptomatic anemia due to MDS associated with a deletion 5q. Mantle Cell Lymphoma (MCL): Diagnosis of MCL. Follicular Lymphoma (FL): Diagnosis of FL that has been previously treated. Used in combination with a rituximab product. Marginal Zone Lymphoma (MZL): Diagnosis of MZL that has been previously treated. Used in combination with a rituximab product.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

REZLIDHIA

Products Affected

- Rezlidhia

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of acute myeloid leukemia (AML). Disease is relapsed or refractory. Positive for a susceptible isocitrate dehydrogenase-1 (IDH1) mutation (e.g., R132C, R132H, R132G, R132S, R132L).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

RINVOQ

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid arthritis (RA) (init): Diagnosis (Dx) of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine. Psoriatic arthritis (PsA) (init): Dx of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Ankylosing spondylitis (AS) (init): Dx of active AS. Non-radiographic axial spondyloarthritis (NRAS) (init): Dx of active NRAS. Patient has signs of inflammation. Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, certolizumab pegol). AS, NRAS (init): Minimum duration of a one-month TF/C/I to one nonsteroidal anti-inflammatory drug (NSAID) (eg, ibuprofen, naproxen) at maximally tolerated doses. Atopic dermatitis (AD) (init): Dx of moderate to severe AD. One of the following: a) Involvement of at least 10% body surface area (BSA), or b) SCORing Atopic Dermatitis (SCORAD) index value of at least 25. Trial and failure of a minimum 30-day supply (14-day supply for topical corticosteroids), contraindication, or intolerance to at least one of the following: a) Medium or higher potency topical corticosteroid, b) Pimecrolimus cream, c) Tacrolimus ointment, or d) Eucrisa ointment. One of the following: 1) Trial and failure of a minimum 12-week supply of at least one systemic drug product for the treatment of AD (examples include, but are not limited to, Adbry, Dupixent, etc.), OR 2) Patient has a contraindication, intolerance, or treatment is inadvisable with both of the following FDA-approved AD therapies: Adbry and Dupixent. Not used in combination with other JAK inhibitors, biologic immunomodulators, or other immunosuppressants (eg, azathioprine, cyclosporine).
Age Restrictions	AD (initial): Patient is 12 years of age or older.
Prescriber Restrictions	RA, PJIA, AS, NRAS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. AD (initial): Prescribed by or in consultation with a dermatologist or allergist/immunologist. CD, UC (initial): Prescribed by or in consultation with a gastroenterologist.

Coverage Duration	RA, PJIA, PsA, AS, NRAS, AD, CD, UC (initial): 6 months, (reauth): Plan year.
Other Criteria	<p>Polyarticular juvenile idiopathic arthritis (PJIA) (init): Dx of active PJIA. Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide. RA, PJIA, PsA, AS (init): Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, etanercept, adalimumab). RA, PJIA, PsA, AS, NRAS (init, reauth): Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (eg, azathioprine, cyclosporine). Crohn's disease (CD) (init): Dx of moderately to severely active CD. One of the following: frequent diarrhea and abdominal pain, at least 10% weight loss, complications (eg, obstruction, fever, abdominal mass), abnormal lab values (eg, CRP), OR CD Activity Index (CDAI) greater than 220. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, azathioprine, corticosteroid (eg, prednisone), methotrexate. Ulcerative colitis (UC) (init): Dx of moderately to severely active UC. One of the following: greater than 6 stools per day, frequent blood in the stools, frequent urgency, presence of ulcers, abnormal lab values (eg, hemoglobin, ESR, CRP), OR dependent on, or refractory to, corticosteroids. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate (eg, mesalamine, olsalazine, sulfasalazine), azathioprine, or corticosteroids (eg, prednisone). CD/UC (init): Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, adalimumab). Not used in combination with other JAK inhibitors, biological therapies for CD/UC, or potent immunosuppressants (eg, azathioprine, cyclosporine). All indications (reauth): Patient demonstrates positive clinical response to therapy. AD (reauth): Not used in combination with other JAK inhibitors, biologic immunomodulators, or other immunosuppressants (eg, azathioprine, cyclosporine). CD/UC (reauth): Not used in combination with other JAK inhibitors, biological therapies for CD/UC, or potent immunosuppressants (eg, azathioprine, cyclosporine).</p>

ROFLUMILAST

Products Affected

- Roflumilast

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD) (initial): Diagnosis of COPD. History of COPD exacerbations which required the use of systemic corticosteroids, antibiotics, or hospital admission. Trial and failure, intolerance, or contraindication to two prior therapies for COPD (e.g., Combivent, Spiriva).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, reauth: plan year
Other Criteria	COPD (reauth): Patient demonstrates positive clinical response to therapy.

ROZLYTREK

Products Affected

- Rozlytrek

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of metastatic non-small cell lung cancer (NSCLC). Patient has ROS1 rearrangement positive tumor(s). Solid Tumors: Patient has solid tumors with a neurotrophic tyrosine receptor kinase (NTRK) gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1, TPR-NTRK1, etc.). Disease is without a known acquired resistance mutation (e.g., TRKA G595R, TRKA G667C or TRKC G623R substitutions). Disease is one of the following: metastatic or unresectable (including cases where surgical resection is likely to result in severe morbidity).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

RUBRACA

Products Affected

- Rubraca

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Ovarian cancer: Diagnosis of ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. Both of the following: 1) Disease is recurrent, and 2) Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin). Presence of deleterious BRCA-mutation (germline and/or somatic). Prostate cancer: Diagnosis of metastatic castration-resistant prostate cancer. Presence of deleterious BRCA mutation. Patient has received previous treatment with both of the following: 1) Androgen receptor-directed therapy [e.g., Erleada (apalutamide), Xtandi (enzalutamide), Zytiga (abiraterone)], AND 2) A taxane-based chemotherapy [e.g., docetaxel, Jevtana (cabazitaxel)]. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog [e.g., Lupron (leuprolide), Zoladex (goserelin)], OR 2) Patient received bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

RYBELSUS

Products Affected

- Rybelsus

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

RYDAPT

Products Affected

- Rydapt

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of acute myeloid leukemia (AML), AML is FMS-like tyrosine kinase 3 (FLT3) mutation-positive, Rydapt will be used in combination with standard induction and consolidation therapy. Aggressive Systemic Mastocytosis (ASM), Systemic Mastocytosis with Associated Hematological Neoplasm (SM-AHN), Mast Cell Leukemia (MCL): Diagnosis of one of the following: aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

SCSEMBLIX

Products Affected

- Scemblix

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of chronic myelogenous/myeloid leukemia (CML). Disease is Philadelphia chromosome-positive (Ph+). Disease is in chronic phase. One of the following: 1) Patient has been previously treated with two or more alternative tyrosine kinase inhibitors (TKI) [e.g., Bosulif (bosutinib), imatinib, Sprycel (dasatinib), Tassigna (nilotinib), Iclusig (ponatinib)] and prescribed medication dose will not exceed 80 mg per day, OR 2) Disease is T315I mutation positive and prescribed medication dose will not exceed 400 mg per day.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

SHINGRIX

Products Affected

- Shingrix

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of herpes zoster (shingles). One of the following: A) Age greater than or equal to 50 years OR B) Both of the following: 1) Age 18 to 49 years and 2) Patient is or will be at increased risk of herpes zoster due to immunodeficiency or immunosuppression caused by known disease or therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months (2 injections per lifetime)
Other Criteria	N/A

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cushing's disease: Diagnosis of endogenous Cushing's disease (i.e, hypercortisolism is not a result of chronic administration of high dose glucocorticoids). Either pituitary surgery has not been curative for the patient OR patient is not a candidate for pituitary surgery.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

SILDENAFIL

Products Affected

- Sildenafil Citrate TABS 20MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH: Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: plan year
Other Criteria	N/A

SIRTURO

Products Affected

- Sirturo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of pulmonary multidrug resistant tuberculosis (MDR-TB), adverse reactions or resistance to standard drugs used to treat MDR-TB, and one of the following: Sirturo is being used in combination with at least 3 other medications to which the patient's MDR-TB isolate has been shown to be susceptible in vitro, or if in vitro testing results are unavailable Sirturo is being used in combination with at least 4 other medications to which the patient's MDR-TB isolate is likely to be susceptible.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	24 weeks
Other Criteria	N/A

SKYCLARYS

Products Affected

- Skyclarys

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of Friedreich's ataxia confirmed via genetic testing demonstrating mutation in the FXN gene. Patient has a Modified Friedreich's Ataxia Rating Scale (mFARS) score of greater than or equal to 20 and less than or equal to 80. Patient has a B-type natriuretic peptide value less than or equal to 200 pg/mL.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with one of the following: Neurologist, Neurogeneticist, or Physiatrist (Physical Medicine and Rehabilitation Specialist).
Coverage Duration	Initial, Reauth: Plan Year.
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy.

SKYRIZI

Products Affected

- Skyrizi INJ 150MG/ML, 180MG/1.2ML, 360MG/2.4ML
- Skyrizi Pen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Plaque Psoriasis (initial): Diagnosis of chronic moderate to severe plaque psoriasis. One of the following: at least 3% body surface area involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Psoriatic arthritis (PsA) (initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Crohn's disease (CD) (Initial): Diagnosis of moderately to severely active CD. Will be used as a maintenance dose following the intravenous induction doses.
Age Restrictions	N/A
Prescriber Restrictions	Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. CD (Initial): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	All uses (initial): 6 months. All uses (reauth): plan year.
Other Criteria	All indications (Reauth): Patient demonstrates positive clinical response to therapy.

SOMAVERT

Products Affected

- Somavert

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acromegaly (Initial): Diagnosis of acromegaly by one of the following: serum growth hormone (GH) level greater than 1 ng/mL after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis, or elevated serum IGF-1 levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis. Inadequate response to one of the following: surgery, radiotherapy, or dopamine agonist (e.g., bromocriptine, cabergoline) therapy or not a candidate for surgery, radiotherapy or dopamine agonist (eg, bromocriptine, cabergoline) therapy. One of the following: 1) Inadequate response, contraindication, or intolerance to a somatostatin analog (e.g., octreotide, lanreotide) or 2) Clinical rationale provided for preferred treatment with pegvisomant (e.g., comorbid diabetes mellitus is present with acromegaly, IGF-1 levels greater than 900 ng/mL). Acromegaly (Reauth): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Acromegaly (Initial, Reauth): plan year
Other Criteria	N/A

SORAFENIB

Products Affected

- Sorafenib Tosylate TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal cell carcinoma (RCC): Both of the following: 1) Diagnosis of RCC, and 2) One of the following: a) Disease has relapsed, or b) Both of the following: i) medically or surgically unresectable tumor, and ii) diagnosis of Stage IV disease. Hepatocellular carcinoma (HCC): Diagnosis of HCC. One of the following: patient has metastatic disease, or patient has extensive liver tumor burden, or patient is inoperable by performance status or comorbidity (local disease or local disease with minimal extrahepatic disease only), or both of the following: a) patient is not a transplant candidate, and b) disease is unresectable. Differentiated thyroid carcinoma (DTC): Diagnosis of one of the following: follicular carcinoma, Hurthle cell carcinoma, or papillary carcinoma. One of the following: metastatic disease, unresectable recurrent disease, or persistent locoregional disease. One of the following: patient has symptomatic disease or patient has progressive disease. Disease is refractory to radioactive iodine (RAI) treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

SOTYKTU

Products Affected

- Sotyktu

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Plaque Psoriasis (initial): Diagnosis of moderate to severe plaque psoriasis. One of the following: at least 3% body surface area (BSA) involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Not used in combination with other potent immunosuppressants (eg, azathioprine, cyclosporine, biologic disease-modifying antirheumatic drugs [DMARDs]).
Age Restrictions	N/A
Prescriber Restrictions	Plaque Psoriasis (initial): Prescribed by or in consultation with a dermatologist
Coverage Duration	Plaque Psoriasis (initial): 6 months. Plaque Psoriasis (reauth): plan year.
Other Criteria	Plaque psoriasis (reauth): Patient demonstrates positive clinical response to therapy. Not used in combination with other potent immunosuppressants (eg, azathioprine, cyclosporine, biologic DMARDs).

SPRYCEL

Products Affected

- Sprycel

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Myeloid Leukemia (CML): Diagnosis (dx) of Philadelphia chromosome-positive/BCR ABL-positive chronic myeloid leukemia (Ph+/BCR ABL+ CML). Acute Lymphoblastic Leukemia (ALL): Diagnosis of Philadelphia chromosome-positive/BCR ABL-positive acute lymphoblastic leukemia (Ph+/BCR ABL+ ALL).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

STELARA

Products Affected

- Stelara INJ 45MG/0.5ML, 90MG/ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Plaque psoriasis (Initial - 45mg/0.5mL): Diagnosis of moderate to severe plaque psoriasis. Plaque psoriasis (Initial - 90mg/1mL): Diagnosis of moderate to severe plaque psoriasis. Patient's weight is greater than 100 kg (220 lbs). Plaque psoriasis (Initial regardless of dose): One of the following: at least 3% body surface area involvement, severe scalp psoriasis, OR palmoplantar (ie, palms, soles), facial, or genital involvement. Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies: corticosteroids (eg, betamethasone, clobetasol), vitamin D analogs (eg, calcitriol, calcipotriene), tazarotene, or calcineurin inhibitors (eg, tacrolimus, pimecrolimus). Psoriatic arthritis (PsA) (Initial - 45mg/0.5mL): Diagnosis of active PsA. PsA (Initial - 90mg/1mL): Diagnosis of active PsA. Patient's weight is greater than 100 kg (220 lbs). Diagnosis of co-existent moderate to severe psoriasis. PsA (Initial regardless of dose): One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Crohn's disease (CD) (initial): Diagnosis of moderately to severely active Crohn's disease. Will be used as a maintenance dose following the intravenous induction dose.</p>
Age Restrictions	N/A
Prescriber Restrictions	Plaque psoriasis (Initial): Prescribed by or in consultation with a dermatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. CD and UC (initial): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	All indications (initial): 6 months. All indications (reauth): plan year.
Other Criteria	Ulcerative colitis (UC) (initial): Diagnosis of moderately to severely active UC. Will be used as a maintenance dose following the intravenous induction dose. All indications (Reauth): Patient demonstrates positive clinical response to therapy.

STIVARGA

Products Affected

- Stivarga

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Colorectal Cancer (CRC): Diagnosis of advanced or metastatic colorectal cancer. Trial and failure, contraindication, or intolerance to treatment with all the following: oxaliplatin-based chemotherapy, irinotecan-based chemotherapy, fluoropyrimidine-based chemotherapy, and anti-VEGF therapy-based chemotherapy. One of the following: 1) Tumor is RAS mutant-type OR 2) Tumor is RAS wild-type and trial and failure, contraindication, or intolerance to anti-EGFR therapy. Gastrointestinal stromal tumor (GIST): Diagnosis of progressive, locally advanced, unresectable or metastatic GIST. One of the following: 1) First-line therapy as a single agent for succinate dehydrogenase (SDH) deficient GIST with gross residual disease (R2 resection) or 2) Trial and failure, contraindication, or intolerance to imatinib mesylate and sunitinib malate. Hepatocellular Carcinoma (HCC): Diagnosis of HCC. Used as subsequent-line therapy for disease progression.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

SUNITINIB

Products Affected

- Sunitinib Malate

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Gastrointestinal stromal tumor (GIST): Diagnosis of GIST. One of the following: 1) History of disease progression on, contraindication, or intolerance to Gleevec (imatinib), Stivarga (regorafenib), or standard dose Qinlock (ripretinib) or 2) Succinate dehydrogenase (SDH)-deficient GIST. Renal Cell Carcinoma (RCC): Diagnosis of RCC. One of the following: 1) Disease has relapsed, 2) Both of the following: used in adjuvant setting and patient has a high risk of recurrence following nephrectomy, or 3) Disease is advanced. Islet Cell Tumors/Pancreatic Neuroendocrine Tumors (pNET): Diagnosis of islet cell tumors/progressive pNET.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

SYMPAZAN

Products Affected

- Sympazan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Lennox-Gastaut syndrome: Diagnosis of Lennox-Gastaut syndrome. Used for the adjunctive treatment of seizures associated with Lennox-Gastaut syndrome. Dravet syndrome: Diagnosis of seizures associated with Dravet syndrome (DS). Used in combination with Diacomit.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TABRECTA

Products Affected

- Tabrecta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of non-small cell lung cancer (NSCLC). One of the following: a) Presence of mesenchymal-epithelial transition (MET) exon 14 skipping positive tumors or b) High level MET amplification in lung cancer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TADALAFIL (BPH)

Products Affected

- Tadalafil TABS 2.5MG, 5MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for the treatment of erectile dysfunction only.
Required Medical Information	Benign prostatic hyperplasia (BPH): Diagnosis of BPH. Male Gender.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	BPH: plan year
Other Criteria	BPH: Trial and failure, contraindication, or intolerance to two formulary alpha blockers (e.g., tamsulosin, alfuzosin). 2.5mg strength: Patient has renal insufficiency.

TADALAFIL (PAH)

Products Affected

- Tadalafil TABS 20MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH): Diagnosis of PAH. PAH is symptomatic. One of the following: A) Diagnosis of PAH was confirmed by right heart catheterization or B) Patient is currently on any therapy for the diagnosis of PAH.
Age Restrictions	N/A
Prescriber Restrictions	PAH: Prescribed by or in consultation with a pulmonologist or cardiologist.
Coverage Duration	PAH: plan year
Other Criteria	N/A

TAFAMIDIS

Products Affected

- Vyndaqel

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM) (initial): Diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM). One of the following: 1) Patient has a transthyretin (TTR) mutation (e.g., V122I), 2) Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits, OR 3) All of the following: i) echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis, ii) scintigraphy scan suggestive of cardiac TTR amyloidosis, and iii) absence of light-chain amyloidosis. One of the following: 1) History of heart failure (HF), with at least one prior hospitalization for HF, OR 2) Presence of clinical signs and symptoms of HF (e.g., dyspnea, edema). Documentation of one of the following: a) Patient has New York Heart Association (NYHA) Functional Class I or II heart failure, or b) Patient has New York Heart Association (NYHA) Functional Class III heart failure and patient's cardiopulmonary functional status allows patient to ambulate 100 meters or greater in six minutes or less.
Age Restrictions	N/A
Prescriber Restrictions	ATTR-CM (initial, reauth): Prescribed by or in consultation with a cardiologist
Coverage Duration	ATTR-CM (initial, reauth): Plan year
Other Criteria	ATTR-CM (reauth): Patient demonstrates positive clinical response to therapy. Patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure.

TAFINLAR

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable melanoma or metastatic melanoma AND cancer is BRAFV600 mutant type. Adjuvant Treatment for Melanoma: Diagnosis of melanoma. Cancer is BRAF V600E or V600K mutant type. Involvement of lymph nodes following complete resection. Used as adjunctive therapy. Medication is used in combination with Mekinist (trametinib). Non-small Cell Lung Cancer (NSCLC): All of the following: diagnosis of metastatic non-small cell lung cancer AND cancer is BRAF V600E mutant type AND medication is used in combination with Mekinist (trametinib). Anaplastic Thyroid Cancer (ATC): Diagnosis of anaplastic thyroid cancer. One of the following: 1) Disease is one of the following: metastatic, locally advanced, or unresectable OR 2) Prescribed as adjuvant therapy following resection. Cancer is BRAF V600E mutant type. Medication is used in combination with Mekinist (trametinib). Solid tumors: Presence of solid tumor. Disease is unresectable or metastatic. Patient has progressed on or following prior systemic treatment (e.g., carboplatin, 5-fluorouracil, paclitaxel). Cancer is BRAF V600E mutant type. Medication is used in combination with Mekinist (trametinib). Low-grade glioma: Diagnosis of low-grade glioma. Cancer is BRAF V600E mutant type. Medication is used in combination with Mekinist (trametinib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TAGRISSO

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): One of the following: A) All of the following: Diagnosis of NSCLC. Disease is one of the following: 1) advanced, 2) recurrent, or 3)metastatic. One of the following: 1) Used as first-line therapy AND One of the following: a) Tumors are positive for epidermal growth factor receptor (EGFR) exon 19 deletions, or b) Tumors are positive for EGFR exon 21 L858R mutations, or c) Disease is sensitizing EGFR mutation positive, OR 2) Tumors are positive for EGFR T790M mutation AND Trial and failure, contraindication, or intolerance to at least one prior EGFR tyrosine kinase inhibitor (TKI) therapy [e.g., Iressa (gefitinib), Tarceva (erlotinib), Gilotrif (afatinib)]. OR B) All of the following: Diagnosis of NSCLC. One of the following: 1) Tumors are positive for epidermal growth factor receptor (EGFR) exon 19 deletions, OR 2) Tumors are positive for EGFR exon 21 L858R mutations. Both of the following: 1) Patient is receiving as adjuvant therapy, AND 2) Patient has had a complete surgical resection of the primary NSCLC tumor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TALZENNA

Products Affected

- Talzenna

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) locally advanced or b) metastatic. Presence of deleterious or suspected deleterious germline BRCA-mutations as detected by the FDA-approved companion diagnostic for Talzenna. Prostate cancer: Diagnosis of prostate cancer. Disease is HRR gene-mutated. Disease is metastatic castration-resistant. Taken in combination with Xtandi (enzalutamide). Used in combination with a gonadotropin-releasing hormone (GnRH) analog [e.g., Lupron (leuprolide), Zoladex (goserelin), Trelstar (triptorelin), Vantas (histrelin), Firmagon (degarelix)] OR Patient has had bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TASIGNA

Products Affected

- Tasigna

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Myeloid Leukemia (CML): Diagnosis of Philadelphia chromosome-positive/BCR ABL-positive chronic myeloid leukemia (Ph+/BCR ABL+ CML).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TASIMELTEON CAPSULE

Products Affected

- Tasimelteon

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-24-hour sleep-wake disorder: Diagnosis of non-24-hour sleep-wake disorder (also known as free-running disorder, free-running or non-entrained type circadian rhythm sleep disorder, or hypnnychthemeral syndrome). Smith-Magenis Syndrome (SMS): Diagnosis of Smith-Magenis Syndrome (SMS). Patient is experiencing nighttime sleep disturbances (i.e., difficulty falling asleep, frequent nighttime waking and early waking).
Age Restrictions	SMS: 16 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Non-24, SMS: 6 months
Other Criteria	Non-24: Patient is totally blind (has no light perception).

TAZAROTENE

Products Affected

- Tazarotene CREA 0.1%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	All indications: Excluded if treatment for cosmetic purposes.
Required Medical Information	Acne vulgaris: Diagnosis of acne vulgaris (i.e., acne). Psoriasis: Diagnosis of psoriasis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

TAZVERIK

Products Affected

- Tazverik

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Epithelioid sarcoma: Diagnosis of epithelioid sarcoma. Disease is one of the following: metastatic or locally advanced. Patient is not eligible for complete resection. Follicular lymphoma: Diagnosis of follicular lymphoma. Subsequent therapy for one of the following: a) EZH2 mutation positive relapsed/refractory disease after two prior therapies, OR b) EZH2 wild-type or unknown relapsed/refractory disease and no satisfactory alternative treatment options.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TEPMETKO

Products Affected

- Tepmetko

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is one of the following: recurrent, advanced, or metastatic. Tumor is MET exon 14 skipping mutation positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TERIPARATIDE

Products Affected

- Forteo INJ 600MCG/2.4ML
- Teriparatide INJ 620MCG/2.48ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Postmenopausal osteoporosis or osteopenia or men with primary or hypogonadal osteoporosis or osteopenia (initial): Diagnosis of one of the following: a) postmenopausal osteoporosis or osteopenia or b) primary or hypogonadal osteoporosis or osteopenia. One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions. Glucocorticoid-Induced Osteoporosis: See Other Criteria section.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	All indications (initial, reauth): plan year.
Other Criteria	Glucocorticoid-Induced Osteoporosis (initial): Diagnosis of glucocorticoid-induced osteoporosis. History of prednisone or its equivalent at a dose greater than or equal to 5mg/day for greater than or equal to 3 months. One of the following: 1) BMD T-score less than or equal to -2.5 based on BMD measurements from lumbar spine, femoral neck, total hip, or radius (one-third radius site), or 2) One of the following FRAX 10-year probabilities: a) Major osteoporotic fracture at 20% or

more in the U.S., or the country-specific threshold in other countries or regions, or b) Hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions, 3) History of one of the following fractures resulting from minimal trauma: vertebral compression fx, fx of the hip, fx of the distal radius, fx of the pelvis, or fx of the proximal humerus, or 4) One of the following: a) glucocorticoid dosing of at least 30 mg per day, or b) cumulative glucocorticoid dosing of at least 5 grams per year. TF/C/I to one bisphosphonate (e.g., alendronate). All uses (initial, reauth): One of the following: 1) Treatment duration of parathyroid hormones (e.g., teriparatide, Tymlos [abaloparatide]) has not exceeded a total of 24 months during the patient's lifetime, or 2) Patient remains at or has returned to having a high risk for fracture despite a total of 24 months of use of parathyroid hormones (e.g., teriparatide, Tymlos [abaloparatide]).

TETRABENAZINE

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Huntington's Disease: Diagnosis of chorea in patients with Huntington's disease. Tardive dyskinesia: Diagnosis of tardive dyskinesia. One of the following: 1) Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication or 2) Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication. Tourette's syndrome: Patient has tics associated with Tourette's syndrome. Trial and failure, contraindication, or intolerance to Haldol (haloperidol).
Age Restrictions	Tardive dyskinesia: Age greater than or equal to 18 years.
Prescriber Restrictions	Huntington's: Prescribed by or in consultation with a neurologist. Tardive dyskinesia, Tourette's: Prescribed by or in consultation with a neurologist or psychiatrist.
Coverage Duration	Plan year.
Other Criteria	N/A

THALOMID

Products Affected

- Thalomid

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Erythema Nodosum Leprosum (ENL): Diagnosis (Dx) of moderate to severe ENL. One of the following: used for acute treatment OR used as maintenance therapy for prevention & suppression of cutaneous manifestations of ENL recurrence. Multiple Myeloma (MM): Dx of multiple myeloma.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TIBSOVO

Products Affected

- Tibsovo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Relapsed or refractory Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is relapsed or refractory. AML is isocitrate dehydrogenase-1 (IDH1) mutation-positive. Newly-Diagnosed AML: Diagnosis of newly-diagnosed AML. AML is isocitrate dehydrogenase-1 (IDH1) mutation-positive. One of the following: 1) Patient is greater than or equal to 75 years old OR 2) Patient has comorbidities that preclude the use of intensive induction chemotherapy OR 3) Patient is greater than or equal to 60 years old and not a candidate for or declines intensive induction therapy OR 4) Patient is greater than or equal to 60 years old and receiving post-induction therapy following response to previous lower intensity therapy. Cholangiocarcinoma: Diagnosis of cholangiocarcinoma. Disease is locally advanced, unresectable, or metastatic. Cholangiocarcinoma is IDH1 mutation-positive. Disease has progressed on or after systemic treatment. Myelodysplastic Syndromes (MDS): Diagnosis of MDS. Disease is one of the following: relapsed or refractory. MDS is IDH1 mutation-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TOBI PODHALER

Products Affected

- Tobi Podhaler

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

TOPICAL RETINOIDS

Products Affected

- Tretinoin CREA
- Tretinoin GEL 0.01%, 0.025%
- Tretinoin Microsphere GEL 0.04%, 0.1%

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	All indications: Excluded if treatment for cosmetic purposes.
Required Medical Information	Acne vulgaris: Diagnosis of acne vulgaris (i.e., acne).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

TORPENZ

Products Affected

- Torpenz

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced Neuroendocrine Tumors (NET): Diagnosis (Dx) of neuroendocrine tumors of pancreatic origin, gastrointestinal origin, lung origin, or thymic origin. Disease is progressive. Disease is unresectable, locally advanced or metastatic. Advanced Renal Cell Carcinoma/Kidney Cancer: Dx of advanced renal cell cancer/kidney cancer. Disease is one of the following: (1) relapsed or (2) stage IV disease. Renal angiomyolipoma with tuberous sclerosis complex (TSC): Dx of renal angiomyolipoma and TSC, not requiring immediate surgery. Subependymal Giant Cell Astrocytoma (SEGA) with tuberous sclerosis (TS): Dx of SEGA associated with TS. Patient is not a candidate for curative surgical resection. Breast Cancer: Dx of recurrent or metastatic breast cancer. One of the following: Disease is hormone receptor positive (HR+) [i.e., estrogen-receptor-positive (ER+) or progesterone-receptor-positive (PR+)] OR both of the following: disease is HR- and disease has clinical characteristics that predict a HR+ tumor. Disease is HER2-negative. One of the following: Patient is a postmenopausal woman, patient is a premenopausal woman being treated with ovarian ablation/suppression, or patient is male. One of the following: A) Both of the following: a) one of the following: 1) Disease progressed while on or within 12 months of non-steroidal aromatase inhibitor [e.g., Arimidex (anastrozole), Femara (letrozole)] therapy or 2) patient was treated with tamoxifen at any time AND b) Used in combination with Aromasin (exemestane) OR B) Used in combination with Fulvestrant or Tamoxifen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	All uses: Approve for continuation of prior therapy if within the past 120 days.

TRIENTINE

Products Affected

- Trientine Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration). Trial and failure, contraindication, or intolerance to a penicillamine product (e.g., Depen, Cuprimine)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

TRULICITY

Products Affected

- Trulicity

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Excluded if used for weight loss only.
Required Medical Information	Diabetes Mellitus (DM): Submission of medical records (e.g., chart notes) confirming diagnosis of type 2 DM.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	N/A

TRUMENBA

Products Affected

- Trumenba

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for prevention of Neisseria meningitidis serogroup B. Patient has not received or will not receive more than two Bexsero or Trumenba vaccines in the previous 12 months. One of the following: 1) Age lesser than or equal to 25 years, or 2) Patient is in a high risk group as defined by the Meningococcal Vaccination: Recommendations of the Advisory Committee on Immunization Practices, United States, 2020.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year (2 vaccinations per year)
Other Criteria	N/A

TRUQAP

Products Affected

- Truqap TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of breast cancer. Disease is one of the following: locally advanced or metastatic. Will be taken in combination with fulvestrant. Disease is hormone receptor (HR)-positive. Disease is human epidermal growth factor receptor 2 (HER2)-negative. Patient has one or more PIK3CA/AKT1/PTEN-alterations. One of the following: A) Following progression on at least one endocrine-based regimen in the metastatic setting (e.g., anastrozole, letrozole, exemestane, tamoxifen, etc.) OR B) Recurrence on or within 12 months of completing adjuvant therapy (e.g., chemotherapy).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TUKYSA

Products Affected

- Tukysa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: Diagnosis of breast cancer. Disease is one of the following: a) advanced unresectable or b) metastatic. Disease is human epidermal growth factor receptor 2 (HER2)-positive. Used in combination with trastuzumab and capecitabine. Patient has been previously treated with an anti-HER2-based regimen (e.g., trastuzumab, pertuzumab, ado-trastuzumab emtansine) in the metastatic setting. Colorectal cancer: Diagnosis of colorectal cancer (HER2-amplified and RAS and BRAF wild-type). Disease is HER2-positive. Disease is one of the following: a) advanced, b) unresectable, c) metastatic. One of the following: a) patient has previously been treated with one of the following regimens: i) fluoropyrimidine-based chemotherapy, ii) oxaliplatin-based chemotherapy, iii) irinotecan-based chemotherapy or b) patient is not appropriate for intensive therapy. Used in combination with trastuzumab.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TURALIO

Products Affected

- Turalio CAPS 125MG

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Tenosynovial Giant Cell Tumor (TGCT): Diagnosis of TGCT. Patient is symptomatic. Patient is not a candidate for surgery due to worsening functional limitation or severe morbidity with surgical removal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

TYENNE SC

Products Affected

- Tyenne INJ 162MG/0.9ML

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Rheumatoid Arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. One of the following: 1) Trial and failure, contraindication, or intolerance (TF/C/I) to two of the following: Enbrel (etanercept), one formulary adalimumab product, Rinvoq (upadacitinib), or Xeljanz/Xeljanz XR (tofacitinib), OR 2) for continuation of prior therapy if within the past 120 days. Systemic Juvenile Idiopathic Arthritis (SJIA) (Initial): Diagnosis of active SJIA. TF/C/I to one of the following conventional therapies at maximally tolerated doses: a) minimum duration of a one month trial of a nonsteroidal anti-inflammatory drug (NSAID) (eg, ibuprofen, naproxen), b) minimum duration of a 3-month trial of methotrexate, or c) minimum duration of a 2-week trial of a systemic glucocorticoid (eg, prednisone). Polyarticular Juvenile Idiopathic Arthritis (PJIA) (Initial): Diagnosis of active PJIA. One of the following: 1) TF/C/I to two of the following: Enbrel (etanercept), one formulary adalimumab product, Rinvoq/Rinvoq LQ, Xeljanz (tofacitinib), OR 2) for continuation of prior therapy if within the past 120 days. Giant Cell Arteritis (GCA) (Initial): Diagnosis of GCA. TF/C/I to a glucocorticoid (eg, prednisone).
Age Restrictions	N/A
Prescriber Restrictions	RA, SJIA, PJIA, GCA (Initial): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	RA, SJIA, PJIA, GCA: 6 mo (init), plan year (reauth).
Other Criteria	RA, SJIA, PJIA, GCA (reauth): Patient demonstrates positive clinical response to therapy.

TYMLOS

Products Affected

- Tymlos

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>One of the following diagnoses: 1) Postmenopausal osteoporosis or osteopenia, OR 2) Primary or hypogonadal osteoporosis or osteopenia.</p> <p>One of the following: Set I) Both of the following: A) Bone mineral density (BMD) T-score of -2.5 or lower in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) trial and failure, contraindication, or intolerance (TF/C/I) to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]), or Set II) Both of the following: A) BMD T-score between -1.0 and -2.5 in the lumbar spine, femoral neck, total hip, or radius (one-third radius site) AND B) One of the following: 1) history of low-trauma fracture of the hip, spine, proximal humerus, pelvis, or distal forearm, or 2) both of the following: i) TF/C/I to one osteoporosis treatment (e.g., alendronate, risedronate, zoledronic acid, Prolia [denosumab]) and ii) one of the following FRAX (Fracture Risk Assessment Tool) 10-year probabilities: a) major osteoporotic fracture at 20% or more in the U.S., or the country-specific threshold in other countries or regions, or b) hip fracture at 3% or more in the U.S., or the country-specific threshold in other countries or regions.</p> <p>Treatment duration of parathyroid hormones (e.g., teriparatide, Tymlos [abaloparatide]) has not exceeded a total of 24 months during the patient's lifetime.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year (up to 24 months per lifetime)
Other Criteria	N/A

UBRELVY

Products Affected

- Ubrelvy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of migraine with or without aura. Will be used for the acute treatment of migraine. Trial and failure or intolerance to one triptan (e.g., eletriptan, rizatriptan, sumatriptan) or a contraindication to all triptans. Medication will not be used in combination with another CGRP inhibitor for the acute treatment of migraines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, Reauth: Plan year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy. Will not be used for preventive treatment of migraine. Medication will not be used in combination with another CGRP inhibitor for the acute treatment of migraines.

VALCHLOR

Products Affected

- Valchlor

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Mycosis fungoides-type cutaneous T-cell lymphoma (MF-CTCL): Both of the following: 1) diagnosis of Stage IA MF-CTCL OR diagnosis of Stage IB MF-CTCL AND 2) patient has received at least one prior skin-directed therapy (e.g., topical corticosteroids, bexarotene topical gel [Targretin topical gel], etc.).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VALTOCO

Products Affected

- Valtoco 10 Mg Dose
- Valtoco 15 Mg Dose
- Valtoco 20 Mg Dose
- Valtoco 5 Mg Dose

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Epilepsy: Diagnosis of epilepsy. Frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VANDETANIB

Products Affected

- Caprelsa

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Thyroid Cancer: Diagnosis of unresectable locally advanced or metastatic medullary thyroid cancer
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VANFLYTA

Products Affected

- Vanflyta

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Diagnosis of AML. Disease is FLT3 internal tandem duplication (ITD) positive. Vanflyta will be used in combination with standard cytarabine and anthracycline induction and cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VAXCHORA

Products Affected

- Vaxchora

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Vaccine is being used for active immunization against disease caused by <i>Vibrio cholerae</i> serogroup O1. Patient is traveling to a cholera-affected area. Age 2 through 64 years.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

VENCLEXTA

Products Affected

- Venclexta
- Venclexta Starting Pack

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL): Diagnosis of CLL or SLL. Acute Myeloid Leukemia (AML): One of the following: 1) Diagnosis of newly diagnosed AML. Used in combination with azacitidine, or decitabine, or low-dose cytarabine. One of the following: age 60 years or older OR comorbidities that preclude use of intensive induction chemotherapy. 2) Diagnosis of relapsed/refractory acute myeloid leukemia (AML). Relapse is greater than or equal to 12 months from most recent disease remission. Venclexta therapy to be given in combination with the patients previous initial successful induction regimen (e.g., azacitidine, decitabine, low-dose cytarabine, etc.)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VEOZAH

Products Affected

- Veozah

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of moderate to severe vasomotor symptoms due to menopause.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial, Reauth: 6 months
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy.

VERQUVO

Products Affected

- Verquvo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Heart Failure (CHF) (initial): Diagnosis of CHF. Patient has an ejection fraction less than 45 percent. Patient has New York Heart Association (NYHA) Class II, III, or IV symptoms. One of the following: A) Patient was hospitalized for heart failure within the last 6 months, or B) Patient used outpatient intravenous diuretics (e.g., bumetanide, furosemide) for heart failure within the last 3 months. Trial and failure, contraindication, or intolerance to two of the following at a maximally tolerated dose: A) One of the following: 1) Angiotensin converting enzyme (ACE) inhibitor (e.g., captopril, enalapril), 2) Angiotensin II receptor blocker (ARB) (e.g., candesartan, valsartan), or 3) Angiotensin receptor-neprilysin inhibitor (ARNI) [e.g., Entresto (sacubitril and valsartan)], B) One of the following: 1) bisoprolol, 2) carvedilol, or 3) metoprolol succinate extended release, C) Sodium-glucose co-transporter 2 (SGLT2) inhibitor [e.g., Jardiance (empagliflozin), Farxiga (dapagliflozin), Xigduo XR (dapagliflozin and metformin)], or D) Mineralocorticoid receptor antagonist (MRA) [e.g., eplerenone, spironolactone].
Age Restrictions	N/A
Prescriber Restrictions	CHF (initial): Prescribed by or in consultation with a cardiologist.
Coverage Duration	CHF (initial, reauth): plan year
Other Criteria	CHF (reauth): Patient demonstrates positive clinical response to therapy.

VERZENIO

Products Affected

- Verzenio

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced, Recurrent, or Metastatic Breast Cancer: Diagnosis of advanced, recurrent, or metastatic breast cancer. Disease is hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative. One of the following: a) used in combination with an aromatase inhibitor (e.g., anastrozole, letrozole, exemestane), OR b) used in combination with Faslodex (fulvestrant) OR c) used as monotherapy and disease has progressed following endocrine therapy and patient has already received at least one prior chemotherapy regimen. Early Breast Cancer: Diagnosis of early breast cancer at high risk of recurrence. Disease is hormone receptor (HR)-positive. Disease is human epidermal growth factor receptor 2 (HER2)-negative. Used in combination with one of the following endocrine therapies: 1) tamoxifen or 2) aromatase inhibitor (e.g., anastrozole, letrozole, exemestane).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VIGABATRIN

Products Affected

- Vigabatrin
- Vigadrone
- Vigpoder

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Complex Partial Seizures (CPS): For use as adjunctive therapy. Infantile Spasms (IS): Diagnosis of infantile spasms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days. CPS: Trial and failure, contraindication, or intolerance (TF/C/I) to two formulary anticonvulsants [eg, Lamictal (lamotrigine), Depakene (valproic acid), Dilantin (phenytoin)].

VIGAFYDE

Products Affected

- Vigafyde

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of infantile spasms. Trial and failure, or intolerance to generic vigabatrin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VITRAKVI

Products Affected

- Vitrakvi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Presence of solid tumors (e.g., salivary gland, soft tissue sarcoma, infantile fibrosarcoma, thyroid cancer, lung, melanoma, colon, etc.). Disease is positive for neurotrophic receptor tyrosine kinase (NTRK) gene fusion (e.g. ETV6-NTRK3, TPM3-NTRK1, LMNA-NTRK1, etc.). Disease is without a known acquired resistance mutation [e.g., TRKA G595R, G623R, G696A, F617L]. Disease is one of the following: metastatic or unresectable. One of the following: Disease has progressed on previous treatment (e.g., surgery, radiotherapy, or systemic therapy) OR Disease has no satisfactory alternative treatments.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VIZIMPRO

Products Affected

- Vizimpro

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small cell lung cancer (NSCLC): Diagnosis of NSCLC. Disease is recurrent, advanced, or metastatic. Disease is positive for one of the following epidermal growth factor receptor (EGFR) mutations: a) exon 19 deletion, b) exon 21 L858R substitution, c) S768I, d) L861Q, or e) G719X.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VONJO

Products Affected

- Vonjo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient has been diagnosed with one of the following: a) Primary myelofibrosis, b) Post-polycythemia vera myelofibrosis, OR c) Post-essential thrombocythemia myelofibrosis. One of the following: 1) Patient has a platelet count below $50 \times 10^9/L$ or Both of the following: 1) Patient has a platelet count greater than or equal to $50 \times 10^9/L$ and 2) History of no response or loss of response to one prior JAK inhibitor (e.g., Jakafi or Inrebic).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VORICONAZOLE INJECTION

Products Affected

- Voriconazole INJ

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Invasive aspergillosis: Diagnosis of invasive aspergillosis (IA). Candidemia: Diagnosis of candidemia. One of the following: (1) patient is non-neutropenic or (2) infection is located in skin, abdomen, kidney, bladder wall, or wounds. Esophageal Candidiasis: Diagnosis of esophageal candidiasis. Mycosis: Diagnosis of fungal infection caused by <i>Scedosporium apiospermum</i> (asexual form of <i>Pseudallescheria boydii</i>) or <i>Fusarium</i> spp. including <i>Fusarium solani</i> . For fusariosis: Patient is intolerant of, or refractory to, other therapy (e.g., liposomal amphotericin B, amphotericin B lipid complex).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 weeks
Other Criteria	N/A

VOSEVI

Products Affected

- Vosevi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Criteria will be applied consistent with current AASLD/IDSA guideline. All patients: Diagnosis of chronic hepatitis C, patient is without decompensated liver disease (defined as Child-Pugh Class B or C), and patient is not receiving Vosevi in combination with another HCV direct acting antiviral agent [e.g., Harvoni, Zepatier].
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with one of the following: Hepatologist, Gastroenterologist, Infectious disease specialist, HIV specialist certified through the American Academy of HIV Medicine.
Coverage Duration	12 to 24 weeks. Criteria will be applied consistent with current AASLD/IDSA guideline.
Other Criteria	N/A

VOTRIENT

Products Affected

- Pazopanib Hydrochloride

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Renal Cell Carcinoma (RCC): Diagnosis of RCC. Soft tissue sarcoma (STS): Diagnosis of advanced STS.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

VOWST

Products Affected

- Vowst

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of recurrent clostridioides difficile infection (CDI) as defined by both of the following: 1) Presence of diarrhea defined as a passage of 3 or more loose bowel movements within a 24-hour period for two consecutive days, and 2) A positive stool test for C.difficile toxin or toxigenic C.difficile. Patient has a history of two or more recurrent episodes of CDI within 12 months. All of the following: 1) Patient has completed at least 10 consecutive days of one of the following antibiotic therapies 2-4 days prior to initiating Vowst: oral vancomycin or Difucid (fidaxomicin), 2) Patient has completed the recommended course of magnesium citrate the day before and at least 8 hours prior to initiating Vowst, and 3) Previous episode of CDI is under control (e.g., less than 3 unformed/loose [i.e., Bristol Stool Scale type 6-7] stools/day for 2 consecutive days).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist or infectious disease specialist.
Coverage Duration	14 days
Other Criteria	N/A

WELIREG

Products Affected

- Welireg

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>von Hippel-Lindau (VHL) disease: Diagnosis of von Hippel-Lindau (VHL) disease. Patient requires therapy for one of the following: a) renal cell carcinoma (RCC), b) central nervous system (CNS) hemangioblastoma, or c) pancreatic neuroendocrine tumor (pNET). Patient does not require immediate surgery. Advanced Renal Cell Carcinoma (RCC): Diagnosis of advanced renal cell carcinoma. Disease has progressed after treatment with both of the following: a) Programmed death receptor 1 (PD-1) or programmed death ligand 1 (PD-L1) checkpoint inhibitor [e.g., Keytruda (pembrolizumab), Opdivo (nivolumab), Jemperli (dostarlimab)] and b) Vascular endothelial growth factor tyrosine kinase inhibitor (VEGF-TKI) [e.g., Inlyta (axitinib), Votrient (pazopanib), Cabometyx (cabozantinib)].</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

XALKORI

Products Affected

- Xalkori

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Non-small Cell Lung Cancer (NSCLC): Diagnosis of advanced, metastatic or recurrent NSCLC. One of the following: a) Tumor is anaplastic lymphoma kinase (ALK)-positive, b) Tumor is ROS1-positive c) Tumor is positive for mesenchymal-epithelial transition (MET) amplification, OR d) Tumor is positive for MET exon 14 skipping mutation. Anaplastic Large Cell Lymphoma (ALCL): Diagnosis of systemic ALCL. Disease is relapsed or refractory. Tumor is anaplastic lymphoma kinase (ALK)-positive. Inflammatory Myofibroblastic Tumor (IMT): Diagnosis of IMT. Tumor is anaplastic lymphoma kinase (ALK)-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

XATMEP

Products Affected

- Xatmep

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute lymphoblastic leukemia (ALL): Diagnosis of acute lymphoblastic leukemia (ALL). Polyarticular juvenile idiopathic arthritis (pJIA) (initial): Diagnosis of active polyarticular juvenile idiopathic arthritis. Trial and failure, contraindication, or intolerance to one nonsteroidal anti-inflammatory drug (NSAID) (e.g., diclofenac, ibuprofen, meloxicam, naproxen).
Age Restrictions	ALL: Patient is 18 years of age or younger. pJIA (initial): Patient is 18 years of age or younger.
Prescriber Restrictions	pJIA (initial): Prescribed by or in consultation with a rheumatologist.
Coverage Duration	ALL: plan year. pJIA (initial, reauth): plan year
Other Criteria	ALL: Approve for continuation of prior therapy if within the past 120 days. pJIA (reauth): Patient demonstrates positive clinical response to therapy

XCOPRI

Products Affected

- Xcopri

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of partial onset seizures.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

XELJANZ

Products Affected

- Xeljanz
- Xeljanz Xr

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Xeljanz tab/Xeljanz XR tab: Rheumatoid arthritis (RA) (Initial): Diagnosis of moderately to severely active RA. Minimum duration of a 3-month trial and failure, contraindication, or intolerance (TF/C/I) to one of the following conventional therapies at maximally tolerated doses: methotrexate, leflunomide, sulfasalazine. Xeljanz tab/Xeljanz XR tab: Psoriatic arthritis (PsA) (Initial): Diagnosis of active PsA. One of the following: actively inflamed joints, dactylitis, enthesitis, axial disease, or active skin and/or nail involvement. Xeljanz tab/Xeljanz XR tab: Ankylosing spondylitis (AS) (Initial): Diagnosis of active AS. Minimum duration of a one-month TF/C/I to one nonsteroidal anti-inflammatory drug (NSAID) (eg, ibuprofen, naproxen) at maximally tolerated doses. RA, PsA, AS (Initial): Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, adalimumab, etanercept). Xeljanz tab/Xeljanz XR tab: Ulcerative colitis (UC) (Initial): Diagnosis of moderately to severely active UC. One of the following: greater than 6 stools per day, frequent blood in the stools, frequent urgency, presence of ulcers, abnormal lab values (eg, hemoglobin, ESR, CRP), OR dependent on, or refractory to, corticosteroids. TF/C/I to one of the following conventional therapies: 6-mercaptopurine, aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine), azathioprine, or corticosteroids (eg, prednisone). Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, adalimumab). Not used in combination with other Janus kinase (JAK) inhibitors, biological therapies for UC, or potent immunosuppressants (eg, azathioprine, cyclosporine).</p>
Age Restrictions	N/A
Prescriber Restrictions	RA, PJIA, AS (initial): Prescribed by or in consultation with a rheumatologist. PsA (initial): Prescribed by or in consultation with a dermatologist or rheumatologist. UC (initial): Prescribed by or in consultation with a gastroenterologist.
Coverage Duration	RA/PJIA/PsA/AS (init): 6 mo. UC (init): 4 mo. RA/PJIA/PsA/AS/UC (reauth): plan year.
Other Criteria	Xeljanz: Polyarticular course juvenile idiopathic arthritis (PJIA) (Initial): Diagnosis of active polyarticular course juvenile idiopathic arthritis.

<p>Minimum duration of a 6-week TF/C/I to one of the following conventional therapies at maximally tolerated doses: leflunomide or methotrexate. Patient has had an inadequate response or intolerance to one or more TNF inhibitors (eg, adalimumab, etanercept). RA, PsA, AS, PJIA (Initial, Reauth): Not used in combination with other JAK inhibitors, biologic disease-modifying antirheumatic drugs (DMARDs), or potent immunosuppressants (eg, azathioprine, cyclosporine). All indications (Reauth): Patient demonstrates positive clinical response to therapy. UC (Reauth): Not used in combination with other JAK inhibitors, biological therapies for UC, or potent immunosuppressants (e.g., azathioprine, cyclosporine).</p>
--

XERMELO

Products Affected

- Xermelo

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Carcinoid syndrome diarrhea (Initial): Diagnosis of carcinoid syndrome diarrhea AND diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g., octreotide, lanreotide) for at least 3 months AND used in combination with SSA therapy.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist
Coverage Duration	Initial: 6 months, Reauth: plan year
Other Criteria	Carcinoid syndrome diarrhea (Reauthorization): Patient demonstrates positive clinical response to therapy.

XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prevention of skeletal-related events in patients with multiple myeloma (MM) and bone metastases from solid tumors (BMST): One of the following: 1) Diagnosis of multiple myeloma OR 2) Diagnosis of solid tumors (eg, breast cancer, kidney cancer, lung cancer, prostate cancer, thyroid cancer) and documented evidence of one or more metastatic bone lesions. Giant cell tumor of bone (GCTB): Diagnosis of giant cell tumor of bone. Tumor is unresectable or surgical resection is likely to result in severe morbidity. Hypercalcemia of malignancy (HCM) (initial): Diagnosis of hypercalcemia of malignancy and refractory to bisphosphonate therapy. Hypercalcemia of malignancy (reauthorization): Patient demonstrates positive clinical response to therapy.
Age Restrictions	N/A
Prescriber Restrictions	GCTB, Hypercalcemia of malignancy (initial): Prescribed by or in consultation with an oncologist
Coverage Duration	MM/BMST: plan year. GCTB: 6 mo. HCM (all): 2 mo.
Other Criteria	Giant cell tumor of bone : Approve for continuation of prior therapy if within the past 120 days.

XIFAXAN

Products Affected

- Xifaxan

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Traveler's Diarrhea (TD): Diagnosis of traveler's diarrhea. Trial and failure, contraindication, or intolerance to one of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin) OR resistance to all of the following: Cipro (ciprofloxacin), Levaquin (levofloxacin), ofloxacin, Zithromax (azithromycin). Prophylaxis of Hepatic Encephalopathy (HE): Used for the prophylaxis of hepatic encephalopathy recurrence. One of the following: 1) Trial and failure, contraindication, or intolerance to lactulose OR 2) Add-on treatment to lactulose. Treatment of HE: Diagnosis of HE. Used for the treatment of HE. One of the following: 1) Trial and failure, contraindication, or intolerance to lactulose OR 2) Add-on treatment to lactulose. Irritable Bowel Syndrome with Diarrhea (Initial): Diagnosis of irritable bowel syndrome with diarrhea (IBS-D). Trial and failure, contraindication or intolerance to an antidiarrheal agent (e.g., loperamide).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	TD: 14 days (one treatment course). HE (Prophylaxis, Tx): plan year. IBS-D (initial/reauth): 2 wks
Other Criteria	IBS-D Reauthorization: Patient experiences IBS-D symptom recurrence.

XOLAIR

Products Affected

- Xolair

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Asthma (Initial): Diagnosis of moderate to severe persistent allergic asthma. Baseline (pre-Xolair treatment) serum total IgE level greater than or equal to 30 IU/mL and less than or equal to 700 IU/mL for patients 12 years of age and older OR greater than or equal to 30 IU/mL and less than or equal to 1300 IU/mL for patients 6 years to less than 12 years of age. Positive skin test or in vitro reactivity to a perennial aeroallergen. One of the following: a) Patient has had two or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months, OR b) Prior asthma-related hospitalization within the past 12 months. Patient is currently being treated with one of the following, unless there is a contraindication or intolerance to these medications: 1) one maximally-dosed combination inhaled corticosteroid/long-acting beta2-agonist [eg, Advair (fluticasone propionate/salmeterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone/vilanterol)] or 2) Both of the following: a) one high-dose inhaled corticosteroid (ICS) [e.g., greater than 500 mcg fluticasone propionate equivalent/day] and b) one additional asthma controller medication {e.g., leukotriene receptor antagonist, long-acting beta2-agonist [eg, Foradil (formoterol fumarate), Serevent (salmeterol xinafoate)], tiotropium}. Chronic Spontaneous Urticaria (CSU) (Previously Chronic Idiopathic Urticaria) (Initial): Diagnosis of CSU (previously chronic idiopathic urticaria). Persistent symptoms (itching and hives) with a second generation H1 antihistamine (e.g., cetirizine, fexofenadine), unless there is a history of contraindication or intolerance to H1 antihistamines. Patient has tried and had an inadequate response or intolerance or contraindication to one of the following additional therapies: H1-antihistamine, Hydroxyzine, H2-antagonist (e.g., famotidine, cimetidine), Leukotriene receptor antagonist (e.g., montelukast).</p>
Age Restrictions	N/A
Prescriber Restrictions	<p>Asthma (Init): Prescribed by or in consultation with a pulmonologist or allergist/immunologist. CSU (Init): Prescribed by or in consultation with an allergist/immunologist, or dermatologist. CRSwNP (Init): Prescribed by or in consultation with an allergist/immunologist, otolaryngologist, or pulmonologist. IgE-Mediated Food Allergy (Init/Reauth): Prescribed by or in consultation with an allergist/immunologist</p>

Coverage Duration	Asthma,Init:6m,Reauth:PlanYr. CSU,Init:3m,reauth:6m. CRSwNP:PlanYr.Allergy, Init:20wk,Reauth:PlanYr
Other Criteria	<p>Chronic Rhinosinusitis with Nasal Polyps (CRSwNP) (Previously Nasal Polyps) (Initial): Diagnosis of CRSwNP (previously nasal polyps). Unless contraindicated, the patient has had an inadequate response to an intranasal corticosteroid (e.g., fluticasone, mometasone). CRSwNP (Reauthorization): Patient demonstrates positive clinical response to therapy. CRSwNP (Initial/Reauth): Used in combination with another agent for chronic rhinosinusitis with nasal polyps. Asthma (Reauthorization): Patient demonstrates positive clinical response to therapy. Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications. CSU (Reauthorization): Patient demonstrates positive clinical response to therapy. IgE-Mediated Food Allergy (Initial): One of the following: A) Both of the following: 1) Diagnosis of IgE Mediated Food Allergy as evidenced by one of the following: a) Positive skin prick test (defined as greater than or equal to 4 mm wheal greater than saline control) to food, b) Positive food specific IgE (greater than or equal to 6 kUA/L), c) Positive oral food challenge, defined as experiencing dose-limiting symptoms at a single dose of less than or equal to 300 mg of food protein, AND 2) Clinical history of IgE Mediated Food Allergy, OR B) Provider attestation that patient has a history of severe allergic response, including anaphylaxis, following exposure to one or more foods. Used in conjunction with food allergen avoidance. Baseline (pre-Xolair treatment) serum total IgE level is greater than or equal to 30 IU/mL and less than or equal to 1850 IU/mL. Dosing is according to serum total IgE levels and body weight. IgE-Mediated Food Allergy (Reauth): Patient demonstrates positive clinical response to therapy. Used in conjunction with food allergen avoidance. Dosing will continue to be based on body weight and pretreatment total IgE serum levels.</p>

XOLREMDI

Products Affected

- Xolremdi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Initial: Diagnosis of WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome. Patient has genotype confirmed variant of CXCR4. Patient has an absolute neutrophil count (ANC) less than or equal to 500 cells/ μ L.
Age Restrictions	N/A
Prescriber Restrictions	Initial: Prescribed by or in consultation with one of the following: dermatologist, immunologist, hematologist, geneticist, or allergist.
Coverage Duration	Initial: 6 months, Reauth: Plan Year
Other Criteria	Reauth: Patient demonstrates positive clinical response to therapy.

XOSPATA

Products Affected

- Xospata

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of acute myeloid leukemia (AML). AML is FMS-like tyrosine kinase 3 (FLT3) mutation-positive. One of the following: a) Used in combination with azacitidine as low-intensity treatment induction when not a candidate for intensive induction therapy, b) Follow-up after induction therapy with response to previous lower intensity therapy with the same regimen, c) Post-allogeneic hematopoietic cell transplantation and in remission, or d) Disease is relapsed or refractory.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

XPOVIO

Products Affected

- Xpovio
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Twice Weekly

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Multiple Myeloma (MM): Diagnosis of multiple myeloma. Patient has received at least one prior therapy (e.g., lenalidomide, bortezomib, daratumumab, pomalidomide). Used in combination with one of the following: bortezomib and dexamethasone, daratumumab and dexamethasone, or carfilzomib and dexamethasone. Relapsed/Refractory Multiple Myeloma (RRMM): Diagnosis of relapsed or refractory multiple myeloma (RRMM). Patient has received at least four prior therapies (e.g., lenalidomide, bortezomib, daratumumab, pomalidomide). Disease is refractory to all of the following: 1) Two proteasome inhibitors (e.g., bortezomib, carfilzomib), 2) Two immunomodulatory agents (e.g., lenalidomide, thalidomide), and 3) An anti-CD38 monoclonal antibody (e.g. daratumumab). Used in combination with dexamethasone. Diffuse large B-cell lymphoma (DLBCL): Diagnosis of one of the following: 1) Relapsed or refractory DLBCL not otherwise specified OR 2) Relapsed or refractory DLBCL arising from follicular lymphoma. Patient has previously received at least two lines of systemic therapy (e.g., CHOP: cyclophosphamide, doxorubicin, vincristine, and prednisone plus rituximab).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

XTANDI

Products Affected

- Xtandi

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prostate cancer (PC): Diagnosis of prostate cancer. One of the following: 1) Disease is castration-resistant, 2) Disease is both of the following: a) Metastatic, and b) Castration-sensitive, or 3) Disease is all of the following: a) Non-metastatic, b) Castration-sensitive, c) Recurrent, and d) High risk of metastasis. One of the following: 1) Used in combination with a gonadotropin-releasing hormone (GnRH) analog [e.g. Lupron (leuprolide), Zoladex (goserelin), Trelstar (triptorelin), Vantas (histrelin), Firmagon (degarelix)] or 2) Patient has had bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZEJULA

Products Affected

- Zejula TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: advanced epithelial ovarian cancer, advanced fallopian tube cancer, or advanced primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to first-line platinum-based chemotherapy (e.g., cisplatin, carboplatin). Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of one of the following: recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer. Used for maintenance treatment in patients who are in a complete or partial response to platinum-based chemotherapy (e.g., cisplatin, carboplatin).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days

ZELBORAF

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Melanoma: Diagnosis of unresectable or metastatic melanoma. Patient is positive for BRAF V600 mutation. Erdheim-Chester Disease: Diagnosis of Erdheim-Chester disease AND Disease is BRAFV600 mutant type (MT).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZOLINZA

Products Affected

- Zolinza

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Cutaneous T-cell lymphoma (CTCL): Diagnosis of CTCL. Patient has progressive, persistent, or recurrent disease on or following two systemic therapies [e.g., Adcetris (brentuximab vedotin), bexarotene, interferon gamma-1b, methotrexate, Poteligeo (mogamulizumab), romidepsin].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZTALMY

Products Affected

- Ztalmy

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD). Patient has a mutation in the CDKL5 gene. Trial and failure, contraindication, or intolerance to two formulary anticonvulsants (e.g., valproic acid, levetiracetam, lamotrigine).
Age Restrictions	Patient is 2 years of age or older.
Prescriber Restrictions	Prescribed by or in consultation with a neurologist.
Coverage Duration	Plan Year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZURZUVAE

Products Affected

- Zurzuvae

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Postpartum Depression (PPD): One of the following: A) Diagnosis of severe PPD or B) Both of the following: a) Diagnosis of mild to moderate PPD, and b) Trial and failure, contraindication, or intolerance to at least one oral SSRI or SNRI (e.g., escitalopram, duloxetine). Onset of symptoms in the third trimester or within 4 weeks of delivery. Prescriber attests that the patient has been counseled and has agreed to adhere to the following: Will follow instructions to not drive or operate machinery until at least 12 hours after taking each dose of Zurzuvae for the duration of the 14-day treatment course and that patients are informed that they may not be able to assess their own driving competence or the degree of driving impairment caused by Zurzuvae.
Age Restrictions	PPD: Patient is 18 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	14 days
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZYDELIG

Products Affected

- Zydelig

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): Diagnosis of CLL or SLL. Disease has relapsed or is refractory.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

ZYKADIA

Products Affected

- Zykadia TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of metastatic, recurrent, or advanced non-small cell lung cancer (NSCLC), tumor is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan year
Other Criteria	Approve for continuation of prior therapy if within the past 120 days.

PART B VERSUS PART D

Products Affected

- Abelcet
- Acetylcysteine INHALATION SOLN
- Acyclovir Sodium INJ 50MG/ML
- Albuterol Sulfate NEBU 0.083%,
0.63MG/3ML, 1.25MG/3ML,
2.5MG/0.5ML
- Amphotericin B INJ
- Amphotericin B Liposome
- Anzemet TABS 50MG
- Arformoterol Tartrate
- Azathioprine TABS 50MG
- Budesonide SUSP
- Calcitriol CAPS
- Calcitriol ORAL SOLN
- Cinacalcet Hydrochloride
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclophosphamide TABS
- Cyclosporine CAPS
- Cyclosporine Modified
- Dextrose 5%
- Dextrose 5%/sodium Chloride 0.9%
- Doxercalciferol CAPS
- Engerix-b
- Envarsus Xr
- Everolimus TABS 0.25MG, 0.5MG,
0.75MG, 1MG
- Formoterol Fumarate NEBU
- Furosemide INJ
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Granisetron Hydrochloride TABS
- Heparin Sodium INJ 1000UNIT/ML
- Hekplisav-b
- Imovax Rabies (h.d.c.v.)
- Intralipid INJ 20GM/100ML,
30GM/100ML
- Ipratropium Bromide INHALATION
SOLN 0.02%
- Ipratropium Bromide/albuterol
Sulfate
- Levalbuterol NEBU

- Levalbuterol Hcl NEBU
0.31MG/3ML, 1.25MG/3ML
- Levalbuterol Hydrochloride NEBU
0.63MG/3ML
- Mycophenolate Mofetil CAPS
- Mycophenolate Mofetil SUSR
- Mycophenolate Mofetil TABS
- Mycophenolic Acid Dr
- Myhibbin
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hydrochloride TABS
- Ondansetron Odt TBDP 4MG, 8MG
- Paricalcitol CAPS
- Pentamidine Isethionate
INHALATION SOLR
- Plenamine INJ 147.4MEQ/L;
2.17GM/100ML; 1.47GM/100ML;
434MG/100ML; 749MG/100ML;
1.04GM/100ML; 894MG/100ML;
749MG/100ML; 1.04GM/100ML;
1.18GM/100ML; 749MG/100ML;
1.04GM/100ML; 894MG/100ML;
592MG/100ML; 749MG/100ML;
250MG/100ML; 39MG/100ML;
960MG/100ML
- Potassium Chloride INJ
10MEQ/100ML, 20MEQ/100ML,
2MEQ/ML, 40MEQ/100ML
- Potassium Chloride/dextrose INJ 5%;
20MEQ/L
- Potassium Chloride/sodium Chloride
INJ 20MEQ/L; 0.45%, 20MEQ/L;
0.9%, 40MEQ/L; 0.9%
- Prehevbrio
- Premasol INJ 52MEQ/L;
1760MG/100ML; 880MG/100ML;
34MEQ/L; 1760MG/100ML;
372MG/100ML; 406MG/100ML;
526MG/100ML; 492MG/100ML;
492MG/100ML; 526MG/100ML;
356MG/100ML; 356MG/100ML;
390MG/100ML; 34MG/100ML;
152MG/100ML
- Prograf PACK

- Prosol
- Pulmozyme SOLN 2.5MG/2.5ML
- Rabavert
- Recombivax Hb
- Sirolimus SOLN
- Sirolimus TABS
- Sodium Chloride INJ 0.9%, 3%, 5%
- Tacrolimus CAPS
- Tobramycin NEBU 300MG/5ML
- Travasol INJ 52MEQ/L;
1760MG/100ML; 880MG/100ML;
34MEQ/L; 1760MG/100ML;
372MG/100ML; 406MG/100ML;
526MG/100ML; 492MG/100ML;
492MG/100ML; 526MG/100ML;
356MG/100ML; 500MG/100ML;
356MG/100ML; 390MG/100ML;
34MG/100ML; 152MG/100ML
- Trophamine INJ 0.54GM/100ML;
1.2GM/100ML; 0.32GM/100ML; 0;
0; 0.5GM/100ML; 0.36GM/100ML;
0.48GM/100ML; 0.82GM/100ML;
1.4GM/100ML; 1.2GM/100ML;
0.34GM/100ML; 0.48GM/100ML;
0.68GM/100ML; 0.38GM/100ML;
5MEQ/L; 0.025GM/100ML;
0.42GM/100ML; 0.2GM/100ML;
0.24GM/100ML; 0.78GM/100ML

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Plans are insured through UnitedHealthcare Insurance Company or one of its affiliated companies, a Medicare Advantage organization with a Medicare contract and a Medicare-approved Part D sponsor. Enrollment in the plan depends on the plan's contract renewal with Medicare.

[<OVEX3386715_000>]

Formulary ID# 00025003

Y0066_130404_093713 CMS Approved